

D52 / TPD52 Antibody (C-Terminus)
Goat Polyclonal Antibody
Catalog # ALS15777**Specification**

D52 / TPD52 Antibody (C-Terminus) - Product Information

Application	WB, IHC-P, E
Primary Accession	P55327
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Calculated MW	24kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A E~~N/A

D52 / TPD52 Antibody (C-Terminus) - Additional Information**Gene ID** 7163**Other Names**

Tumor protein D52, Protein N8, TPD52

Target/Specificity

Human TPD52. This antibody is expected to recognise all three reported isoforms (NP_001020423.1, NP_001020424.1 and NP_005070.1).

Reconstitution & Storage

Store at -20°C. Minimize freezing and thawing.

Precautions

D52 / TPD52 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

D52 / TPD52 Antibody (C-Terminus) - Protein Information**Name** TPD52**Tissue Location**


Isoform 2 is expressed in colon, breast, prostate, pancreas and kidney tumor cell lines. Isoform 2 is expressed at high levels in kidney, prostate, brain, small intestine and pancreas, at moderate levels in placenta and colon, at low levels in lung, liver and heart, and at very low levels in spleen, thymus, peripheral mononuclear blood cells, testis and ovary.

D52 / TPD52 Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

D52 / TPD52 Antibody (C-Terminus) - Images



Antibody (1 ug/ml) staining of Human Lymph Node lysate (35 ug protein in RIPA buffer).

D52 / TPD52 Antibody (C-Terminus) - References

Byrne J.A., et al. Cancer Res. 55:2896-2903(1995).
Chen S.-L., et al. Oncogene 12:741-751(1996).
Wang R., et al. Cancer Res. 64:1589-1594(2004).
Wang R., et al. Biochem. Biophys. Res. Commun. 389:455-460(2009).
Ota T., et al. Nat. Genet. 36:40-45(2004).