

## HSD3B7 Antibody (aa121-170)

Rabbit Polyclonal Antibody Catalog # ALS15901

### **Specification**

## HSD3B7 Antibody (aa121-170) - Product Information

Application IHC, WB Primary Accession Q9H2F3

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 41kDa KDa

### HSD3B7 Antibody (aa121-170) - Additional Information

#### Gene ID 80270

#### **Other Names**

3 beta-hydroxysteroid dehydrogenase type 7, 3 beta-hydroxysteroid dehydrogenase type VII, 3-beta-HSD VII, 3-beta-hydroxy-Delta(5)-C27 steroid oxidoreductase, C(27) 3-beta-HSD, 1.1.1.-, Cholest-5-ene-3-beta, 7-alpha-diol 3-beta-dehydrogenase, 1.1.1.181, HSD3B7

## **Target/Specificity**

HSD3B7 Antibody detects endogenous levels of total HSD3B7 protein.

## **Reconstitution & Storage**

Store at -20°C for up to one year.

### **Precautions**

HSD3B7 Antibody (aa121-170) is for research use only and not for use in diagnostic or therapeutic procedures.

#### HSD3B7 Antibody (aa121-170) - Protein Information

## Name HSD3B7 (HGNC:18324)

### **Function**

The 3-beta-HSD enzymatic system plays a crucial role in the biosynthesis of all classes of hormonal steroids. HSD VII is active against four 7-alpha-hydroxylated sterols. Does not metabolize several different C(19/21) steroids as substrates. Involved in bile acid synthesis (PubMed:<a href="http://www.uniprot.org/citations/11067870" target="\_blank">11067870</a>). Plays a key role in cell positioning and movement in lymphoid tissues by mediating degradation of 7-alpha,25- dihydroxycholesterol (7-alpha,25-OHC): 7-alpha,25-OHC acts as a ligand for the G protein-coupled receptor GPR183/EBI2, a chemotactic receptor for a number of lymphoid cells (By similarity).

#### **Cellular Location**

Endoplasmic reticulum membrane; Multi-pass membrane protein




**Volume** 50 μl

## HSD3B7 Antibody (aa121-170) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## HSD3B7 Antibody (aa121-170) - Images




Anti-HSD3B7 antibody IHC staining of human kidney.



Western blot of extracts from HeLa/MCF-7/COLO cells, using HSD3B7 Antibody.

# HSD3B7 Antibody (aa121-170) - Background

The 3-beta-HSD enzymatic system plays a crucial role in the biosynthesis of all classes of hormonal steroids. HSD VII is active against four 7-alpha-hydroxylated sterols. Does not metabolize several different C(19/21) steroids as substrates. Involved in bile acid synthesis.





# HSD3B7 Antibody (aa121-170) - References

Schwarz M., et al.J. Clin. Invest. 106:1175-1184(2000).
Ota T., et al.Nat. Genet. 36:40-45(2004).
Martin J., et al.Nature 432:988-994(2004).
Mural R.J., et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Cheng J.B., et al.J. Clin. Endocrinol. Metab. 88:1833-1841(2003).