

SPIN / SPIN1 Antibody (aa111-160)
Rabbit Polyclonal Antibody
Catalog # ALS15981**Specification**

SPIN / SPIN1 Antibody (aa111-160) - Product Information

Application	WB
Primary Accession	Q9Y657
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	30kDa KDa

SPIN / SPIN1 Antibody (aa111-160) - Additional Information**Gene ID** 10927**Other Names**

Spindlin-1, Ovarian cancer-related protein, Spindlin1, SPIN1, OCR, SPIN

Target/Specificity

SPIN1 Antibody detects endogenous levels of total SPIN1 protein.

Reconstitution & Storage

Store at -20°C for up to one year.

Precautions

SPIN / SPIN1 Antibody (aa111-160) is for research use only and not for use in diagnostic or therapeutic procedures.

SPIN / SPIN1 Antibody (aa111-160) - Protein Information**Name** SPIN1**Synonyms** OCR, SPIN**Function**

Chromatin reader that specifically recognizes and binds histone H3 both trimethylated at 'Lys-4' and asymmetrically dimethylated at 'Arg-8' (H3K4me3 and H3R8me2a) and acts as an activator of Wnt signaling pathway downstream of PRMT2. In case of cancer, promotes cell cancer proliferation via activation of the Wnt signaling pathway (PubMed:[24589551](http://www.uniprot.org/citations/24589551)). Overexpression induces metaphase arrest and chromosomal instability. Localizes to active rDNA loci and promotes the expression of rRNA genes (PubMed:[21960006](http://www.uniprot.org/citations/21960006)). May play a role in cell- cycle regulation during the transition from gamete to embryo. Involved in oocyte meiotic resumption, a process that takes place before ovulation to resume meiosis of oocytes blocked in prophase I: may act by regulating maternal transcripts to control meiotic resumption.

Cellular Location

Nucleus. Nucleus, nucleolus

Tissue Location

Highly expressed in ovarian cancer tissues.


Volume

50 µl

SPIN / SPIN1 Antibody (aa111-160) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SPIN / SPIN1 Antibody (aa111-160) - Images

Western blot of extracts from HeLa cells, using SPIN1 Antibody.

SPIN / SPIN1 Antibody (aa111-160) - Background

Chromatin reader that specifically recognizes and binds histone H3 both trimethylated at 'Lys-4' and asymmetrically dimethylated at 'Arg-8' (H3K4me3 and H3R8me2a) and acts as an activator of Wnt signaling pathway downstream of PRMT2. In case of cancer, promotes cell cancer proliferation via activation of the Wnt signaling pathway (PubMed:24589551). Overexpression induces metaphase arrest and chromosomal instability. Localizes to active rDNA loci and promotes the expression of rRNA genes (PubMed:21960006). May play a role in cell-cycle regulation during the transition from gamete to embryo. Involved in oocyte meiotic resumption, a process that takes place before ovulation to resume meiosis of oocytes blocked in prophase I: may act by regulating maternal transcripts to control meiotic resumption.

SPIN / SPIN1 Antibody (aa111-160) - References

Peng Y.,et al.Submitted (NOV-1998) to the EMBL/GenBank/DDBJ databases.
Zhang H.L.,et al.Chin. Sci. Bull. 45:909-914(2000).
Gao Y.,et al.Biochem. Biophys. Res. Commun. 335:343-350(2005).
Wiemann S.,et al.Genome Res. 11:422-435(2001).

Kalnine N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.