

**NEFH / NF-H Antibody**  
**Chicken Polyclonal Antibody**  
**Catalog # ALS15985**

### Specification

#### NEFH / NF-H Antibody - Product Information

Application	WB, IHC-P, ICC
Primary Accession	<a href="#">P12036</a>
Reactivity	Human, Bovine
Host	Chicken
Clonality	Polyclonal
Calculated MW	112kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A ICC~~N/A

#### NEFH / NF-H Antibody - Additional Information

##### Gene ID 4744

##### Other Names

Neurofilament heavy polypeptide, NF-H, 200 kDa neurofilament protein, Neurofilament triplet H protein, NEFH, KIAA0845, NFH

##### Reconstitution & Storage

May be stored at 4°C. For long-term storage, aliquot and store at 4°C. Do not freeze. Aliquots are stable for at least 12 months. Sensitive to light.

##### Precautions

NEFH / NF-H Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### NEFH / NF-H Antibody - Protein Information

##### Name NEFH

##### Synonyms KIAA0845, NFH

##### Function

Neurofilaments usually contain three intermediate filament proteins: NEFL, NEFM, and NEFH which are involved in the maintenance of neuronal caliber. NEFH has an important function in mature axons that is not subserved by the two smaller NF proteins. May additionally cooperate with the neuronal intermediate filament proteins PRPH and INA to form neuronal filamentous networks (By similarity).


##### Cellular Location

Cytoplasm, cytoskeleton. Cell projection, axon {ECO:0000250|UniProtKB:P19246}

**Volume**50  $\mu$ l**NEFH / NF-H Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**NEFH / NF-H Antibody - Images**

Anti-NEFH / NF-H antibody IHC staining of human brain, cerebellum.

**NEFH / NF-H Antibody - Background**

Neurofilaments usually contain three intermediate filament proteins: L, M, and H which are involved in the maintenance of neuronal caliber. NF-H has an important function in mature axons that is not subserved by the two smaller NF proteins.

**NEFH / NF-H Antibody - References**

Lees J.F., et al. EMBO J. 7:1947-1955(1988).  
Zhu Y., et al. Beijing Yi Ke Da Xue Xue Bao 31:531-531(1999).  
Nagase T., et al. DNA Res. 5:355-364(1998).  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Dunham I., et al. Nature 402:489-495(1999).