

**FHIT Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS16032****Specification****FHIT Antibody (C-Terminus) - Product Information**

Application	WB, IF, IHC
Primary Accession	P49789
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	17kDa KDa

FHIT Antibody (C-Terminus) - Additional Information**Gene ID** 2272**Other Names**

Bis(5'-adenosyl)-triphosphatase, 3.6.1.29, AP3A hydrolase, AP3Aase, Diadenosine 5', 5'''-P1, P3-triphosphate hydrolase, Dinucleosidetriphosphatase, Fragile histidine triad protein, FHIT

Target/Specificity

FHIT antibody is human and mouse reactive.

Reconstitution & Storage

Long term: -20°C; Short term: +4°C. Avoid repeat freeze-thaw cycles.

Precautions

FHIT Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

FHIT Antibody (C-Terminus) - Protein Information**Name** FHIT**Function**

Possesses dinucleoside triphosphate hydrolase activity (PubMed:12574506, PubMed:15182206, PubMed:8794732, PubMed:9323207, PubMed:9576908, PubMed:9543008). Cleaves P(1)-P(3)-bis(5'-adenosyl) triphosphate (Ap3A) to yield AMP and ADP (PubMed:12574506, PubMed:15182206, PubMed:8794732, PubMed:9323207, PubMed:9576908, PubMed:9543008).

[9576908](http://www.uniprot.org/citations/9576908), PubMed: [9543008](http://www.uniprot.org/citations/9543008)). Can also hydrolyze P(1)-P(4)-bis(5'-adenosyl) tetraphosphate (Ap4A), but has extremely low activity with ATP (PubMed: [8794732](http://www.uniprot.org/citations/8794732)). Exhibits adenylylsulfatase activity, hydrolyzing adenosine 5'-phosphosulfate to yield AMP and sulfate (PubMed: [18694747](http://www.uniprot.org/citations/18694747)). Exhibits adenosine 5'-monophosphoramidase activity, hydrolyzing purine nucleotide phosphoramidates with a single phosphate group such as adenosine 5'-monophosphoramidate (AMP-NH₂) to yield AMP and NH₂ (PubMed: [18694747](http://www.uniprot.org/citations/18694747)). Exhibits adenylylsulfate-ammonia adenylyltransferase, catalyzing the ammonolysis of adenosine 5'-phosphosulfate resulting in the formation of adenosine 5'-phosphoramidate (PubMed: [26181368](http://www.uniprot.org/citations/26181368)). Also catalyzes the ammonolysis of adenosine 5-phosphorofluoridate and diadenosine triphosphate (PubMed: [26181368](http://www.uniprot.org/citations/26181368)). Modulates transcriptional activation by CTNNB1 and thereby contributes to regulate the expression of genes essential for cell proliferation and survival, such as CCND1 and BIRC5 (PubMed: [18077326](http://www.uniprot.org/citations/18077326)). Plays a role in the induction of apoptosis via SRC and AKT1 signaling pathways (PubMed: [16407838](http://www.uniprot.org/citations/16407838)). Inhibits MDM2-mediated proteasomal degradation of p53/TP53 and thereby plays a role in p53/TP53-mediated apoptosis (PubMed: [15313915](http://www.uniprot.org/citations/15313915)). Induction of apoptosis depends on the ability of FHIT to bind P(1)-P(3)-bis(5'-adenosyl) triphosphate or related compounds, but does not require its catalytic activity, it may in part come from the mitochondrial form, which sensitizes the low-affinity Ca(2+) transporters, enhancing mitochondrial calcium uptake (PubMed: [12574506](http://www.uniprot.org/citations/12574506), PubMed: [19622739](http://www.uniprot.org/citations/19622739)). Functions as a tumor suppressor (By similarity).

Cellular Location

Cytoplasm. Mitochondrion. Nucleus

Tissue Location


Low levels expressed in all tissues tested. Phospho-FHIT observed in liver and kidney, but not in brain and lung Phospho-FHIT undetected in all tested human tumor cell lines


FHIT Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FHIT Antibody (C-Terminus) - Images






Western blot analysis of FHIT in HeLa cell lysate with FHIT antibody at (A) 1 and (B) 2 ug/ml.



Immunofluorescence of FHIT in HeLa cells with FHIT antibody at 5 ug/ml.



Anti-FHIT antibody IHC staining of human kidney.

FHIT Antibody (C-Terminus) - Background

Cleaves P(1)-P(3)-bis(5'-adenosyl) triphosphate (Ap3A) to yield AMP and ADP. Can also hydrolyze P(1)-P(4)-bis(5'-adenosyl) tetraphosphate (Ap4A), but has extremely low activity with ATP. Modulates transcriptional activation by CTNNB1 and thereby contributes to regulate the expression of genes essential for cell proliferation and survival, such as CCND1 and BIRC5. Plays a role in the

induction of apoptosis via SRC and AKT1 signaling pathways. Inhibits MDM2-mediated proteasomal degradation of p53/TP53 and thereby plays a role in p53/TP53-mediated apoptosis. Induction of apoptosis depends on the ability of FHIT to bind P(1)-P(3)-bis(5'-adenosyl) triphosphate or related compounds, but does not require its catalytic activity, it may in part come from the mitochondrial form, which sensitizes the low- affinity Ca(2+) transporters, enhancing mitochondrial calcium uptake. Functions as tumor suppressor.

FHIT Antibody (C-Terminus) - References

Ohta M.,et al.Cell 84:587-597(1996).
Druck T.,et al.Cancer Res. 57:504-512(1997).
Corominas R.,et al.Nat. Commun. 5:3650-3650(2014).
Naqvi S.R.A.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).