

**GAPDH Antibody (clone 1D4)**  
**Mouse Monoclonal Antibody**  
**Catalog # ALS16379****Specification****GAPDH Antibody (clone 1D4) - Product Information**

Application	WB, IHC-P, IF
Primary Accession	<a href="#">P04406</a>
Reactivity	Human, Mouse, Rat, Pig
Host	Mouse
Clonality	Monoclonal
Calculated MW	36kDa KDa
Dilution	WB~~1:1000 IHC-P~~N/A IF~~1:50~200

**GAPDH Antibody (clone 1D4) - Additional Information****Gene ID** 2597**Other Names**Glyceraldehyde-3-phosphate dehydrogenase, GAPDH, 1.2.1.12, Peptidyl-cysteine S-nitrosylase  
GAPDH, 2.6.99.-, GAPDH, GAPD**Target/Specificity**

Clone 1D4 is known to react with GAPDH from human, cow, pig, mouse, rat, other mammals, and avian GAPDH.

**Reconstitution & Storage**

Store at -20°C for up to one year.

**Precautions**

GAPDH Antibody (clone 1D4) is for research use only and not for use in diagnostic or therapeutic procedures.

**GAPDH Antibody (clone 1D4) - Protein Information****Name** GAPDH {ECO:0000303|PubMed:2987855, ECO:0000312|HGNC:HGNC:4141}**Function**

Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively (PubMed:<a href="http://www.uniprot.org/citations/11724794" target="\_blank">11724794</a>, PubMed:<a href="http://www.uniprot.org/citations/3170585" target="\_blank">3170585</a>).

Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D- glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate (PubMed:<a href="http://www.uniprot.org/citations/11724794" target="\_blank">11724794</a>, PubMed:<a href="http://www.uniprot.org/citations/3170585" target="\_blank">3170585</a>).

target="\_blank">3170585</a>). Modulates the organization and assembly of the cytoskeleton (By similarity). Facilitates the CHP1- dependent microtubule and membrane associations through its ability to stimulate the binding of CHP1 to microtubules (By similarity). Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes (PubMed:<a href="http://www.uniprot.org/citations/23071094" target="\_blank">23071094</a>). Upon interferon-gamma treatment assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation (PubMed:<a href="http://www.uniprot.org/citations/23071094" target="\_blank">23071094</a>). Also plays a role in innate immunity by promoting TNF-induced NF-kappa-B activation and type I interferon production, via interaction with TRAF2 and TRAF3, respectively (PubMed:<a href="http://www.uniprot.org/citations/23332158" target="\_blank">23332158</a>, PubMed:<a href="http://www.uniprot.org/citations/27387501" target="\_blank">27387501</a>). Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis (By similarity). Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity).

#### Cellular Location

Cytoplasm, cytosol. Nucleus {ECO:0000250|UniProtKB:P04797}. Cytoplasm, perinuclear region. Membrane Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:P04797} Note=Translocates to the nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization signal (By similarity). Postnuclear and Perinuclear regions (PubMed:12829261) {ECO:0000250|UniProtKB:P04797, ECO:0000269|PubMed:12829261}

#### Volume


50 µl

#### GAPDH Antibody (clone 1D4) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### GAPDH Antibody (clone 1D4) - Images



Human Lung: Formalin-Fixed, Paraffin-Embedded (FFPE)

#### **GAPDH Antibody (clone 1D4) - Background**

Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively. Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis. Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC. Modulates the organization and assembly of the cytoskeleton. Facilitates the CHP1-dependent microtubule and membrane associations through its ability to stimulate the binding of CHP1 to microtubules (By similarity). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D-glyceraldehyde 3-phosphate (G3P) into 3-phospho-D- glyceroyl phosphate. Component of the GAIT (gamma interferon- activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes. Upon interferon-gamma treatment assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation.

#### **GAPDH Antibody (clone 1D4) - References**

Hanauer A.,et al.EMBO J. 3:2627-2633(1984).  
Arcari P.,et al.Nucleic Acids Res. 12:9179-9189(1984).  
Tso J.Y.,et al.Nucleic Acids Res. 13:2485-2502(1985).  
Tokunaga K.,et al.Cancer Res. 47:5616-5619(1987).  
Allen R.W.,et al.J. Biol. Chem. 262:649-653(1987).