

15-PGDH / HPGD Antibody (aa205-266)
Rabbit Polyclonal Antibody
Catalog # ALS16382**Specification**

15-PGDH / HPGD Antibody (aa205-266) - Product Information

Application	WB, ICC, IHC
Primary Accession	P15428
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29kDa KDa

15-PGDH / HPGD Antibody (aa205-266) - Additional Information**Gene ID** 3248**Other Names**

15-hydroxyprostaglandin dehydrogenase [NAD(+)], 15-PGDH, 1.1.1.141, Prostaglandin dehydrogenase 1, HPGD, PGDH1

Target/Specificity

Human 15-PGDH / HPGD

Reconstitution & Storage

Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

Precautions

15-PGDH / HPGD Antibody (aa205-266) is for research use only and not for use in diagnostic or therapeutic procedures.

15-PGDH / HPGD Antibody (aa205-266) - Protein Information**Name** HPGD ([HGNC:5154](#))**Synonyms** PGDH1, SDR36C1**Function**

Catalyzes the NAD-dependent dehydrogenation (oxidation) of a broad array of hydroxylated polyunsaturated fatty acids (mainly eicosanoids and docosanoids, including prostaglandins, lipoxins and resolvins), yielding their corresponding keto (oxo) metabolites (PubMed:10837478, PubMed:16757471, PubMed:16828555, PubMed:21916491, PubMed:25586183, PubMed:8086429). Decreases the

levels of the pro- proliferative prostaglandins such as prostaglandin E2 (whose activity is increased in cancer because of an increase in the expression of cyclooxygenase 2) and generates oxo-fatty acid products that can profoundly influence cell function by abrogating pro-inflammatory cytokine expression (PubMed:15574495, PubMed:25586183). Converts resolvins E1, D1 and D2 to their oxo products, which represents a mode of resolvin inactivation. Resolvin E1 plays important roles during the resolution phase of acute inflammation, while resolvins D1 and D2 have a unique role in obesity-induced adipose inflammation (PubMed:16757471, PubMed:22844113).

Cellular Location

Cytoplasm.

Tissue Location


Detected in colon epithelium (at protein level).

15-PGDH / HPGD Antibody (aa205-266) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


15-PGDH / HPGD Antibody (aa205-266) - Images



Sample (30g whole cell lysate). A: Raji . 12% SDS PAGE. 15-PGDH / HPGD antibody diluted at 1:1000



Immunofluorescence of paraformaldehyde-fixed HeLa using 15-PGDH antibody at 1:200 dilution.



Human Small Intestine: Formalin-Fixed, Paraffin-Embedded (FFPE)

15-PGDH / HPGD Antibody (aa205-266) - Background

Prostaglandin inactivation. Contributes to the regulation of events that are under the control of prostaglandin levels. Catalyzes the NAD-dependent dehydrogenation of lipoxin A4 to form 15-oxo-lipoxin A4. Inhibits in vivo proliferation of colon cancer cells.

15-PGDH / HPGD Antibody (aa205-266) - References

Krook M.,et al.Biochemistry 29:738-743(1990).
Ensor C.M.,et al.J. Biol. Chem. 265:14888-14891(1990).
Pichaud F.,et al.Gene 162:319-322(1995).
Delage-Mourroux R.,et al.Gene 188:143-148(1997).
Ota T.,et al.Nat. Genet. 36:40-45(2004).