

BCL3 / BCL-3 Antibody (aa2-13)
Rabbit Polyclonal Antibody
Catalog # ALS16571**Specification**

BCL3 / BCL-3 Antibody (aa2-13) - Product Information

Application	IHC, WB
Primary Accession	P20749
Other Accession	602
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	47584

BCL3 / BCL-3 Antibody (aa2-13) - Additional Information**Gene ID** 602**Other Names**

BCL3, B-cell CLL/lymphoma 3, B-cell leukemia/lymphoma 3, B-cell lymphoma 3 protein, BCL-3, BCL4, D19S37, Proto-oncogene BCL3

Target/Specificity

This Polyclonal antibody is directed against human Bcl 3 protein. The product was purified from serum by protein A chromatography. Expect reactivity with human Bcl 3. Cross-reactivity with Bcl 3 from other sources has not been determined.

Reconstitution & Storage

PBS, pH 7.4, 0.02% sodium azide. Store at -20°C for up to one year.

Precautions

BCL3 / BCL-3 Antibody (aa2-13) is for research use only and not for use in diagnostic or therapeutic procedures.

BCL3 / BCL-3 Antibody (aa2-13) - Protein Information**Name** BCL3**Synonyms** BCL4, D19S37**Function**

Contributes to the regulation of transcriptional activation of NF-kappa-B target genes. In the cytoplasm, inhibits the nuclear translocation of the NF-kappa-B p50 subunit. In the nucleus, acts as transcriptional activator that promotes transcription of NF-kappa-B target genes. Contributes to the regulation of cell proliferation (By similarity).

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, perinuclear region. Note=Ubiquitination via 'Lys-63'- linked ubiquitin chains is required for nuclear accumulation

Volume


50 µl

BCL3 / BCL-3 Antibody (aa2-13) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BCL3 / BCL-3 Antibody (aa2-13) - Images



Human Small Intestine: Formalin-Fixed, Paraffin-Embedded (FFPE)



Lane1: Jurkat cell lysate(20 µg); Lane2: Jurkat cell lysate(40 µg);Bcl3:60kDa

BCL3 / BCL-3 Antibody (aa2-13) - Background

Contributes to the regulation of transcriptional activation of NF-kappa-B target genes. In the cytoplasm, inhibits the nuclear translocation of the NF-kappa-B p50 subunit. In the nucleus, acts as transcriptional activator that promotes transcription of NF-kappa-B target genes. Contributes to the regulation of cell proliferation (By similarity).

BCL3 / BCL-3 Antibody (aa2-13) - References

Ohno H.,et al.Cell 60:991-997(1990).
McKeithan T.W.,et al.Genomics 24:120-126(1994).
Bours V.,et al.Cell 72:729-739(1993).
Heissmeyer V.,et al.EMBO J. 18:4766-4778(1999).
Dechend R.,et al.Oncogene 18:3316-3323(1999).