

PRPF19 / PRP19 Antibody (aa171-220)
Rabbit Polyclonal Antibody
Catalog # ALS16634**Specification****PRPF19 / PRP19 Antibody (aa171-220) - Product Information**

Application	IHC, WB
Primary Accession	O9UMS4
Other Accession	27339
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	55181

PRPF19 / PRP19 Antibody (aa171-220) - Additional Information**Gene ID** 27339**Other Names**

PRPF19, HPSO4, Senescence evasion factor, NMP200, Nuclear matrix protein 200, Pre-mRNA-processing factor 19, PRP19/PSO4 homolog, PRP19, PSO4, SNEV, UBOX4

Target/Specificity

PRPF19 Antibody detects endogenous levels of total PRPF19 protein.

Reconstitution & Storage

PBS (without Mg²⁺, Ca²⁺), pH 7.4, 150 mM sodium chloride, 0.02% sodium azide, 50% glycerol. Store at -20°C for up to one year.

Precautions

PRPF19 / PRP19 Antibody (aa171-220) is for research use only and not for use in diagnostic or therapeutic procedures.

PRPF19 / PRP19 Antibody (aa171-220) - Protein Information**Name** PRPF19 ([HGNC:17896](#))**Function**

Ubiquitin-protein ligase which is a core component of several complexes mainly involved pre-mRNA splicing and DNA repair. Required for pre-mRNA splicing as component of the spliceosome (PubMed: [28502770](http://www.uniprot.org/citations/28502770), PubMed: [28076346](http://www.uniprot.org/citations/28076346), PubMed: [29360106](http://www.uniprot.org/citations/29360106), PubMed: [29301961](http://www.uniprot.org/citations/29301961), PubMed: [30705154](http://www.uniprot.org/citations/30705154)). Core component of the PRP19C/Prp19 complex/NTC/Nineteen complex which is part of the spliceosome and participates in its assembly, its remodeling and is

required for its activity. During assembly of the spliceosome, mediates 'Lys-63'-linked polyubiquitination of the U4 spliceosomal protein PRPF3. Ubiquitination of PRPF3 allows its recognition by the U5 component PRPF8 and stabilizes the U4/U5/U6 tri- snRNP spliceosomal complex (PubMed:20595234). Recruited to RNA polymerase II C-terminal domain (CTD) and the pre-mRNA, it may also couple the transcriptional and spliceosomal machineries (PubMed:21536736). The XAB2 complex, which contains PRPF19, is also involved in pre-mRNA splicing, transcription and transcription-coupled repair (PubMed:17981804). Beside its role in pre-mRNA splicing PRPF19, as part of the PRP19-CDC5L complex, plays a role in the DNA damage response/DDR. It is recruited to the sites of DNA damage by the RPA complex where PRPF19 directly ubiquitinates RPA1 and RPA2. 'Lys-63'-linked polyubiquitination of the RPA complex allows the recruitment of the ATR-ATRIP complex and the activation of ATR, a master regulator of the DNA damage response (PubMed:24332808). May also play a role in DNA double-strand break (DSB) repair by recruiting the repair factor SETMAR to altered DNA (PubMed:18263876). As part of the PSO4 complex may also be involved in the DNA interstrand cross-links/ICLs repair process (PubMed:16223718). In addition, may also mediate 'Lys-48'-linked polyubiquitination of substrates and play a role in proteasomal degradation (PubMed:11435423). May play a role in the biogenesis of lipid droplets (By similarity). May play a role in neural differentiation possibly through its function as part of the spliceosome (By similarity).

Cellular Location

Nucleus. Nucleus, nucleoplasm. Cytoplasm, cytoskeleton, spindle. Cytoplasm. Lipid droplet {ECO:0000250|UniProtKB:Q99KP6}. Note=Nucleoplasmic in interphase cells Irregularly distributed in anaphase cells. In prophase cells, uniformly distributed, but not associated with condensing chromosomes. Found in extrachromosomal regions in metaphase cells. Mainly localized to the mitotic spindle apparatus when chromosomes segregate during anaphase When nuclei reform during late telophase, uniformly distributed in daughter cells and displays no preferred association with decondensing chromatin. Recruited on damaged DNA at sites of double-strand break

Tissue Location

Ubiquitous. Weakly expressed in senescent cells of different tissue origins. Highly expressed in tumor cell lines

Volume


50 µl

PRPF19 / PRP19 Antibody (aa171-220) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PRPF19 / PRP19 Antibody (aa171-220) - Images



Anti-PRPF19 / PRP19 antibody IHC staining of human testis.



Western blot of extracts from COLO/Jurkat cells, using PRPF19 Antibody.

PRPF19 / PRP19 Antibody (aa171-220) - Background

Ubiquitin-protein ligase which is a core component of several complexes mainly involved pre-mRNA splicing and DNA repair. Core component of the PRP19C/Prp19 complex/NTC/Nineteen complex which is part of the spliceosome and participates in its assembly, its remodeling and is required for its activity. During assembly of the spliceosome, mediates 'Lys-63'-linked polyubiquitination of the U4 spliceosomal protein PRPF3. Ubiquitination of PRPF3 allows its recognition by the U5 component PRPF8 and stabilizes the U4/U5/U6 tri-snRNP spliceosomal complex (PubMed:20595234). Recruited to RNA polymerase II C-terminal domain (CTD) and the pre-mRNA, it may also couple the transcriptional and spliceosomal machineries (PubMed:21536736). The XAB2 complex, which contains PRPF19, is also involved in pre- mRNA splicing, transcription and transcription-coupled repair (PubMed:17981804). Beside its role in pre-mRNA splicing PRPF19, as part of the PRP19-CDC5L complex, plays a role in the DNA damage response/DDR. It is recruited to the sites of DNA damage by the RPA complex where PRPF19 directly ubiquitinates RPA1 and RPA2. 'Lys-63'-linked polyubiquitination of the RPA complex allows the recruitment of the ATR-ATRIP complex and the activation of ATR, a master regulator of the DNA damage response (PubMed:24332808). May also play a role in DNA double-strand break (DSB) repair by recruiting the repair factor SETMAR to altered DNA (PubMed:18263876). As part of the PSO4 complex may also be involved in the DNA interstrand cross-links/ICLs repair process (PubMed:16223718). In addition, may also mediate 'Lys-48'-linked polyubiquitination of substrates and play a role in proteasomal degradation (PubMed:11435423). May play a role in the biogenesis of lipid droplets (By similarity). May play a role in neural differentiation possibly through its function as part of the spliceosome (By similarity).

PRPF19 / PRP19 Antibody (aa171-220) - References

Gotzmann J.,et al.Exp. Cell Res. 261:166-179(2000).
Mahajan K.N.,et al.Proc. Natl. Acad. Sci. U.S.A. 100:10746-10751(2003).

Bienvenut W.V.,et al.Submitted (MAR-2009) to UniProtKB.
Bienvenut W.V.,et al.Submitted (JAN-2010) to UniProtKB.
Gerner C.,et al.J. Cell. Biochem. 74:145-151(1999).