

NME2 Antibody (aa91-140)
Rabbit Polyclonal Antibody
Catalog # ALS16689**Specification**

NME2 Antibody (aa91-140) - Product Information

Application	IHC, WB
Primary Accession	P22392
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	17298

NME2 Antibody (aa91-140) - Additional Information**Gene ID** 4831;654364**Other Names**

Nucleoside diphosphate kinase B, NDK B, NDP kinase B, 2.7.4.6, C-myc purine-binding transcription factor PUF, Histidine protein kinase NDKB, 2.7.13.3, nm23-H2, NME2, NM23B

Target/Specificity

NM23 Antibody detects endogenous levels of total NM23 protein.

Reconstitution & Storage

PBS (without Mg²⁺, Ca²⁺), pH 7.4, 150 mM sodium chloride, 0.02% sodium azide, 50% glycerol. Store at -20°C for up to one year.

Precautions

NME2 Antibody (aa91-140) is for research use only and not for use in diagnostic or therapeutic procedures.

NME2 Antibody (aa91-140) - Protein Information**Name** NME2**Synonyms** NM23B**Function**

Major role in the synthesis of nucleoside triphosphates other than ATP. The ATP gamma phosphate is transferred to the NDP beta phosphate via a ping-pong mechanism, using a phosphorylated active-site intermediate (By similarity). Negatively regulates Rho activity by interacting with AKAP13/LBC (PubMed: [15249197](http://www.uniprot.org/citations/15249197)). Acts as a transcriptional activator of the MYC gene; binds DNA non-specifically (PubMed: [8392752](http://www.uniprot.org/citations/8392752)), PubMed: [19435876](http://www.uniprot.org/citations/19435876)). Binds to both single-stranded guanine- and cytosine-rich

strands within the nuclease hypersensitive element (NHE) III(1) region of the MYC gene promoter. Does not bind to duplex NHE III(1) (PubMed:19435876). Has G-quadruplex (G4) DNA-binding activity, which is independent of its nucleotide-binding and kinase activity. Binds both folded and unfolded G4 with similar low nanomolar affinities. Stabilizes folded G4s regardless of whether they are prefolded or not (PubMed:25679041). Exhibits histidine protein kinase activity (PubMed:20946858).

Cellular Location

Cytoplasm. Cell projection, lamellipodium. Cell projection, ruffle. Note=Colocalizes with ITGB1 and ITGB1BP1 at the edge or peripheral ruffles and lamellipodia during the early stages of cell spreading on fibronectin or collagen but not on vitronectin or laminin substrates [Isoform 3]: Cytoplasm. Cytoplasm, perinuclear region. Nucleus

Tissue Location

[Isoform 1]: Ubiquitously expressed.

Volume


50 µl

NME2 Antibody (aa91-140) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NME2 Antibody (aa91-140) - Images



Anti-NME2 / NM23 antibody IHC staining of human spleen.



Western blot of extracts from K562 cells, using NM23 Antibody.