

CXCL3 / GRO Gamma Antibody (aa150-200)
Rabbit Polyclonal Antibody
Catalog # ALS16920**Specification**

CXCL3 / GRO Gamma Antibody (aa150-200) - Product Information

Application	IHC
Primary Accession	P19876
Other Accession	2921
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	11342

CXCL3 / GRO Gamma Antibody (aa150-200) - Additional Information**Gene ID** 2921**Other Names**

CXCL3, C-X-C motif chemokine 3, GRO3 oncogene, Growth-regulated protein gamma, GRO3, GRO-gamma, GRO-gamma(1-73), MGSA gamma, MIP2B, MIP-2b, MIP2-beta, CINC-2b, GROg, SCYB3, Small inducible cytokine b3

Target/Specificity

Human CXCL3

Reconstitution & Storage

PBS, pH 7.2, 0.05% sodium azide. Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Precautions

CXCL3 / GRO Gamma Antibody (aa150-200) is for research use only and not for use in diagnostic or therapeutic procedures.

CXCL3 / GRO Gamma Antibody (aa150-200) - Protein Information**Name** CXCL3**Synonyms** GRO3, GROG, SCYB3**Function**

Ligand for CXCR2 (By similarity). Has chemotactic activity for neutrophils. May play a role in inflammation and exert its effects on endothelial cells in an autocrine fashion. In vitro, the processed form GRO-gamma(5-73) shows a fivefold higher chemotactic activity for neutrophilic granulocytes.

Cellular Location

Secreted.


Volume
50 µl

CXCL3 / GRO Gamma Antibody (aa150-200) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CXCL3 / GRO Gamma Antibody (aa150-200) - Images



Anti-CXCL3 / GRO Gamma antibody IHC staining of human lung.

CXCL3 / GRO Gamma Antibody (aa150-200) - Background

Ligand for CXCR2 (By similarity). Has chemotactic activity for neutrophils. May play a role in inflammation and exert its effects on endothelial cells in an autocrine fashion. In vitro, the processed form GRO-gamma(5-73) shows a fivefold higher chemotactic activity for neutrophilic granulocytes.

CXCL3 / GRO Gamma Antibody (aa150-200) - References

Tekamp-Olson P., et al. J. Exp. Med. 172:911-919(1990).
Haskill S., et al. Proc. Natl. Acad. Sci. U.S.A. 87:7732-7736(1990).
Hillier L.W., et al. Nature 434:724-731(2005).
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Zhang Z., et al. Protein Sci. 13:2819-2824(2004).