

Dipeptidylpeptidase 8 / DPP8 Antibody (aa349-662)
Rabbit Polyclonal Antibody
Catalog # ALS16962**Specification****Dipeptidylpeptidase 8 / DPP8 Antibody (aa349-662) - Product Information**

Application	IHC, WB
Primary Accession	Q6V1X1
Other Accession	54878
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	103358

Dipeptidylpeptidase 8 / DPP8 Antibody (aa349-662) - Additional Information**Gene ID** 54878**Other Names**

DPP8, Dipeptidyl-peptidase 8, DPP8, Dipeptidyl peptidase VIII, Dipeptidylpeptidase 8, DPP8-1, Dipeptidyl peptidase 8, MSTP141, Prolyl dipeptidase DPP8, DP8, DPP VIII

Target/Specificity

Human Dipeptidylpeptidase 8 / DPP8

Reconstitution & Storage

0.1 M Tris-glycine, pH 7.0, 10% glycerol, 0.01% Thimerosal. Keep as concentrated solution. Aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.

Precautions

Dipeptidylpeptidase 8 / DPP8 Antibody (aa349-662) is for research use only and not for use in diagnostic or therapeutic procedures.

Dipeptidylpeptidase 8 / DPP8 Antibody (aa349-662) - Protein Information**Name** DPP8 {ECO:0000303|PubMed:11012666, ECO:0000312|HGNC:HGNC:16490}**Function**

Dipeptidyl peptidase that cleaves off N-terminal dipeptides from proteins having a Pro or Ala residue at position 2 (PubMed:11012666, PubMed:12534281, PubMed:12662155, PubMed:15039077, PubMed:15664838, PubMed:20536396, PubMed:29382749). Acts as a key inhibitor of caspase-1-dependent monocyte and

macrophage pyroptosis in resting cells by preventing activation of NLRP1 and CARD8 (PubMed:27820798, PubMed:29967349, PubMed:32796818). Sequesters the cleaved C-terminal part of NLRP1 and CARD8, which respectively constitute the active part of the NLRP1 and CARD8 inflammasomes, in a ternary complex, thereby preventing their oligomerization and activation (PubMed:34019797, PubMed:33731929, PubMed:33731932). The dipeptidyl peptidase activity is required to suppress NLRP1 and CARD8; however, neither NLRP1 nor CARD8 are bona fide substrates of DPP8, suggesting the existence of substrate(s) required for NLRP1 and CARD8 inhibition (By similarity).

Cellular Location

Cytoplasm

Tissue Location


Ubiquitously expressed, with highest levels in testis, placenta, prostate, muscle and brain

Dipeptidylpeptidase 8 / DPP8 Antibody (aa349-662) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Dipeptidylpeptidase 8 / DPP8 Antibody (aa349-662) - Images



Anti-Dipeptidylpeptidase 8 / DPP8 antibody IHC staining of human testis.



Sample (30 ug of whole cell lysate).



Sample (50 ug of whole cell lysate).

Dipeptidylpeptidase 8 / DPP8 Antibody (aa349-662) - Background

Dipeptidyl peptidase that cleaves off N-terminal dipeptides from proteins having a Pro or Ala residue at position 2. May play a role in T-cell activation and immune function.

Dipeptidylpeptidase 8 / DPP8 Antibody (aa349-662) - References

- Abbott C.A.,et al.Eur. J. Biochem. 267:6140-6150(2000).
- Qi S.Y.,et al.Biochem. J. 373:179-189(2003).
- Sha J.H.,et al.Submitted (JUL-2003) to the EMBL/GenBank/DDBJ databases.
- Ota T.,et al.Nat. Genet. 36:40-45(2004).
- Zhao B.,et al.Submitted (AUG-1999) to the EMBL/GenBank/DDBJ databases.