

TLR9 Antibody (C-Terminus)
Rabbit Polyclonal Antibody
Catalog # ALS17221**Specification**

TLR9 Antibody (C-Terminus) - Product Information

Application	IHC-P, IF, WB
Primary Accession	O9NR96
Other Accession	54106
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	115860

TLR9 Antibody (C-Terminus) - Additional Information**Gene ID** 54106**Other Names**

TLR9, CD289, CD289 antigen, Scri2a, Toll-like receptor 9

Target/Specificity

peptide corresponding to 16 amino acids near the carboxy terminus of human TLR9

Reconstitution & Storage

PBS, 0.02% sodium azide. Short term 4°C, long term aliquot and store at -20°C, avoid freeze thaw cycles. Store undiluted.

Precautions

TLR9 Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

TLR9 Antibody (C-Terminus) - Protein Information**Name** TLR9**Function**

Key component of innate and adaptive immunity. TLRs (Toll-like receptors) control host immune response against pathogens through recognition of molecular patterns specific to microorganisms. TLR9 is a nucleotide-sensing TLR which is activated by unmethylated cytidine-phosphate-guanosine (CpG) dinucleotides (PubMed: [14716310](http://www.uniprot.org/citations/14716310)). Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed: [11564765](http://www.uniprot.org/citations/11564765), PubMed: [17932028](http://www.uniprot.org/citations/17932028)). Controls lymphocyte response to Helicobacter infection (By similarity). Upon CpG stimulation, induces B-cell proliferation, activation, survival and antibody production (PubMed: [23857366](http://www.uniprot.org/citations/23857366))

target="_blank">23857366).

Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q9EQU3}. Early endosome membrane. Lysosome {ECO:0000250|UniProtKB:Q9EQU3} Cytoplasmic vesicle, phagosome {ECO:0000250|UniProtKB:Q9EQU3}. Golgi apparatus membrane. Note=Relocalizes from endoplasmic reticulum to endosome and lysosome upon stimulation with agonist. Exit from the ER requires UNC93B1. Endolysosomal localization is required for proteolytic cleavage and subsequent activation Intracellular localization of the active receptor may prevent from responding to self nucleic acid. {ECO:0000250|UniProtKB:Q9EQU3, ECO:0000269|PubMed:14716310, ECO:0000269|PubMed:38169466}

Tissue Location


Highly expressed in spleen, lymph node, tonsil and peripheral blood leukocytes, especially in plasmacytoid pre-dendritic cells. Levels are much lower in monocytes and CD11c+ immature dendritic cells. Also detected in lung and liver

TLR9 Antibody (C-Terminus) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


TLR9 Antibody (C-Terminus) - Images




Human Kidney: Formalin-Fixed, Paraffin-Embedded (FFPE)



Human Liver: Formalin-Fixed, Paraffin-Embedded (FFPE)



Immunofluorescence of TLR9 in Mouse Spleen cells with TLR9 antibody at 10 ug/ml.



Western blot of TLR9 in mouse spleen cell lysate with TLR9 antibody at (A) 0.5, (B) 1 and (C) 2...

TLR9 Antibody (C-Terminus) - Background

Key component of innate and adaptive immunity. TLRs (Toll-like receptors) control host immune response against pathogens through recognition of molecular patterns specific to microorganisms. TLR9 is a nucleotide-sensing TLR which is activated by unmethylated cytidine-phosphate-guanosine (CpG) dinucleotides. Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Controls lymphocyte response to Helicobacter infection.

TLR9 Antibody (C-Terminus) - References

Du X.,et al.Eur. Cytokine Netw. 11:362-371(2000).
Chuang T.-H.,et al.Eur. Cytokine Netw. 11:372-378(2000).
Hemmi H.,et al.Nature 408:740-745(2000).
Liu Z.,et al.Submitted (SEP-2007) to the EMBL/GenBank/DDBJ databases.
Nakajima T.,et al.Immunogenetics 60:727-735(2008).