

Anti-KCNJ16 / Kir5.1 Antibody (aa376-428)

Rabbit Anti Human Polyclonal Antibody Catalog # ALS17326

Specification

Anti-KCNJ16 / Kir5.1 Antibody (aa376-428) - Product Information

Application WB, IHC-P, IF

Primary Accession Q9NPI9

Predicted Human, Mouse, Rat Host Rabbit Clonality Polyclonal

Calculated MW 47949

Dilution WB~~1:1000 IHC-P~~N/A IF~~1:50~200

Anti-KCNJ16 / Kir5.1 Antibody (aa376-428) - Additional Information

Gene ID 3773

Alias Symbol KCNJ16
Other Names

KCNJ16, BIR9, KIR5.1

Target/Specificity Human KCNJ16 / Kir5.1

Reconstitution & Storage

PBS, pH 7.2, 0.05% sodium azide Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Precautions

Anti-KCNJ16 / Kir5.1 Antibody (aa376-428) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-KCNJ16 / Kir5.1 Antibody (aa376-428) - Protein Information

Name KCNJ16

Function

Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. KCNJ16 may be involved in the regulation of fluid and pH balance. In the kidney, together with KCNJ10, mediates basolateral K(+) recycling in distal tubules; this process is critical for Na(+) reabsorption at the tubules (PubMed:<ah href="http://www.uniprot.org/citations/24561201" target="_blank">24561201).



Cellular Location

Membrane; Multi- pass membrane protein. Basolateral cell membrane. Note=In kidney distal convoluted tubules, located in the basolateral membrane in the presence of KCNJ10

Tissue Location


Widely expressed, with highest levels in adult and fetal kidney (at protein level). In the kidney, expressed in the proximal and distal convoluted tubules, but not in glomeruli nor collecting ducts.

Anti-KCNJ16 / Kir5.1 Antibody (aa376-428) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-KCNJ16 / Kir5.1 Antibody (aa376-428) - Images



Human Brain, Cortex: Formalin-Fixed, Paraffin-Embedded (FFPE)