

Anti-CD32C Antibody (C-Terminus)
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17424**Specification**

Anti-CD32C Antibody (C-Terminus) - Product Information

Application	WB, IHC-P
Primary Accession	P31995
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	35578

Anti-CD32C Antibody (C-Terminus) - Additional Information**Gene ID** 9103**Alias Symbol** FCGR2C**Other Names**

FCGR2C, CD32, CD32 antigen, CDw32, Fc gamma receptor IIC, FCG2, Fc-gamma-RIIc, FcRII-c, FcRIIC, Igg fc receptor ii-c, Fc-gamma rii-c, HFcRII-C, CD32C, IGFR2

Target/Specificity

Recognizes endogenous levels of CD32c protein.

Reconstitution & Storage

PBS, pH 7.3, 0.01% sodium azide, 30% glycerol. Store at -20°C. Aliquot to avoid freeze/thaw cycles.

Precautions

Anti-CD32C Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-CD32C Antibody (C-Terminus) - Protein Information**Name** FCGR2C**Synonyms** CD32, FCG2, IGFR2**Function**

Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells.

Cellular Location

[Isoform IIC4]: Cytoplasm. [Isoform IIC2]: Cell membrane; Single-pass type I membrane protein

Tissue Location

Isoform IIC1 is detected in monocytes, macrophages, polymorphonuclear cells and natural killer cells

Anti-CD32C Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CD32C Antibody (C-Terminus) - Images