

Anti-EAR2 / NR2F6 Antibody (N-Terminus)
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS17485**Specification**

Anti-EAR2 / NR2F6 Antibody (N-Terminus) - Product Information

Application	IHC-P
Primary Accession	P10588
Predicted	Human, Monkey
Host	Rabbit
Clonality	Polyclonal
Calculated MW	42979

Anti-EAR2 / NR2F6 Antibody (N-Terminus) - Additional Information**Gene ID** 2063**Alias Symbol** NR2F6**Other Names**

NR2F6, Erba related 2, ERBAL2, EAR-2, EAR2, ERBA-related gene-2, V-erbA-related protein 2

Target/Specificity

Human NR2F6. BLAST analysis of the peptide immunogen showed no homology with other human proteins.

Reconstitution & Storage

Immunoaffinity purified

Precautions

Anti-EAR2 / NR2F6 Antibody (N-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-EAR2 / NR2F6 Antibody (N-Terminus) - Protein Information**Name** NR2F6**Synonyms** EAR2, ERBAL2**Function**

Transcription factor predominantly involved in transcriptional repression. Binds to promoter/enhancer response elements that contain the imperfect 5'-AGGTCA-3' direct or inverted repeats with various spacings which are also recognized by other nuclear hormone receptors. Involved in modulation of hormonal responses. Represses transcriptional activity of the lutropin-choriogonadotropic hormone receptor/LHCGR gene, the renin/REN gene and the oxytocin-neurophysin/OXT gene. Represses the triiodothyronine- dependent and -independent transcriptional activity of the thyroid hormone receptor gene in a cell type-specific manner. The corepressing function towards thyroid hormone receptor beta/THRB involves at least in part the inhibition of THRB binding to triiodothyronine response elements (TREs) by NR2F6. Inhibits NFATC

transcription factor DNA binding and subsequently its transcriptional activity. Acts as transcriptional repressor of IL-17 expression in Th-17 differentiated CD4(+) T cells and may be involved in induction and/or maintenance of peripheral immunological tolerance and autoimmunity. Involved in development of forebrain circadian clock; is required early in the development of the locus coeruleus (LC).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00407, ECO:0000269|PubMed:10644740, ECO:0000269|PubMed:18701084}

Tissue Location

Expressed in heart, placenta, liver, skeletal muscle, kidney and pancreas.

Anti-EAR2 / NR2F6 Antibody (N-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-EAR2 / NR2F6 Antibody (N-Terminus) - Images