

**Anti-RIPK1 / RIP Antibody (C-Terminus)**  
**Rabbit Anti Human Polyclonal Antibody**  
**Catalog # ALS17651****Specification****Anti-RIPK1 / RIP Antibody (C-Terminus) - Product Information**

Application	IHC-P
Primary Accession	<a href="#">Q13546</a>
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	75931
Dilution	IHC-P~~N/A

**Anti-RIPK1 / RIP Antibody (C-Terminus) - Additional Information****Gene ID 8737**Alias Symbol **RIPK1****Other Names**

RIPK1, Cell death protein RIP, Receptor interacting protein, RIP1, RIP, RIP-1, Protein kinase rip, Receptor-interacting protein 1, Rinp

**Reconstitution & Storage**

Immunoaffinity purified

**Precautions**

Anti-RIPK1 / RIP Antibody (C-Terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-RIPK1 / RIP Antibody (C-Terminus) - Protein Information****Name** RIPK1 ([HGNC:10019](#))**Function**

Serine-threonine kinase which is a key regulator of TNF- mediated apoptosis, necroptosis and inflammatory pathways (PubMed:<a href="http://www.uniprot.org/citations/17703191" target="\_blank">17703191</a>, PubMed:<a href="http://www.uniprot.org/citations/24144979" target="\_blank">24144979</a>, PubMed:<a href="http://www.uniprot.org/citations/31827280" target="\_blank">31827280</a>, PubMed:<a href="http://www.uniprot.org/citations/31827281" target="\_blank">31827281</a>, PubMed:<a href="http://www.uniprot.org/citations/32657447" target="\_blank">32657447</a>, PubMed:<a href="http://www.uniprot.org/citations/35831301" target="\_blank">35831301</a>). Exhibits kinase activity-dependent functions that regulate cell death and kinase-independent scaffold functions regulating inflammatory signaling and cell survival (PubMed:<a href="http://www.uniprot.org/citations/11101870" target="\_blank">11101870</a>, PubMed:<a href="http://www.uniprot.org/citations/19524512" target="\_blank">19524512</a>, PubMed:<a href="http://www.uniprot.org/citations/19524513" target="\_blank">19524513</a>, PubMed:<a href="http://www.uniprot.org/citations/29440439" target="\_blank">29440439</a>).

target="\_blank">29440439</a>, PubMed:<a href="http://www.uniprot.org/citations/30988283" target="\_blank">30988283</a>). Has kinase-independent scaffold functions: upon binding of TNF to TNFR1, RIPK1 is recruited to the TNF-R1 signaling complex (TNF-RSC also known as complex I) where it acts as a scaffold protein promoting cell survival, in part, by activating the canonical NF-kappa-B pathway (By similarity). Kinase activity is essential to regulate necroptosis and apoptosis, two parallel forms of cell death: upon activation of its protein kinase activity, regulates assembly of two death-inducing complexes, namely complex IIa (RIPK1-FADD-CASP8), which drives apoptosis, and the complex IIb (RIPK1-RIPK3-MLKL), which drives necroptosis (By similarity). RIPK1 is required to limit CASP8- dependent TNFR1-induced apoptosis (By similarity). In normal conditions, RIPK1 acts as an inhibitor of RIPK3-dependent necroptosis, a process mediated by RIPK3 component of complex IIb, which catalyzes phosphorylation of MLKL upon induction by ZBP1 (PubMed:<a href="http://www.uniprot.org/citations/19524512" target="\_blank">19524512</a>, PubMed:<a href="http://www.uniprot.org/citations/19524513" target="\_blank">19524513</a>, PubMed:<a href="http://www.uniprot.org/citations/29440439" target="\_blank">29440439</a>, PubMed:<a href="http://www.uniprot.org/citations/30988283" target="\_blank">30988283</a>). Inhibits RIPK3- mediated necroptosis via FADD-mediated recruitment of CASP8, which cleaves RIPK1 and limits TNF-induced necroptosis (PubMed:<a href="http://www.uniprot.org/citations/19524512" target="\_blank">19524512</a>, PubMed:<a href="http://www.uniprot.org/citations/19524513" target="\_blank">19524513</a>, PubMed:<a href="http://www.uniprot.org/citations/29440439" target="\_blank">29440439</a>, PubMed:<a href="http://www.uniprot.org/citations/30988283" target="\_blank">30988283</a>). Required to inhibit apoptosis and necroptosis during embryonic development: acts by preventing the interaction of TRADD with FADD thereby limiting aberrant activation of CASP8 (By similarity). In addition to apoptosis and necroptosis, also involved in inflammatory response by promoting transcriptional production of pro-inflammatory cytokines, such as interleukin-6 (IL6) (PubMed:<a href="http://www.uniprot.org/citations/31827280" target="\_blank">31827280</a>, PubMed:<a href="http://www.uniprot.org/citations/31827281" target="\_blank">31827281</a>). Phosphorylates RIPK3: RIPK1 and RIPK3 undergo reciprocal auto- and trans- phosphorylation (PubMed:<a href="http://www.uniprot.org/citations/19524513" target="\_blank">19524513</a>). Phosphorylates DAB2IP at 'Ser-728' in a TNF-alpha-dependent manner, and thereby activates the MAP3K5-JNK apoptotic cascade (PubMed:<a href="http://www.uniprot.org/citations/15310755" target="\_blank">15310755</a>, PubMed:<a href="http://www.uniprot.org/citations/17389591" target="\_blank">17389591</a>). Required for ZBP1-induced NF-kappa-B activation in response to DNA damage (By similarity).

#### Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q60855}. Cell membrane {ECO:0000250|UniProtKB:Q9ZUF4}

#### Anti-RIPK1 / RIP Antibody (C-Terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-RIPK1 / RIP Antibody (C-Terminus) - Images