

# Anti-CFI / Complement Factor I Antibody (Internal)

Rabbit Anti Human Polyclonal Antibody Catalog # ALS17749

## **Specification**

# Anti-CFI / Complement Factor I Antibody (Internal) - Product Information

Application WB, IHC-P
Primary Accession P05156
Predicted Human
Host Rabbit
Clonality Polyclonal
Calculated MW 65750

Dilution WB~~1:1000 IHC-P~~N/A

# Anti-CFI / Complement Factor I Antibody (Internal) - Additional Information

**Gene ID 3426** 

**Other Names** 

Alias Symbol

CFI, AHUS3, C3BINA, C3b-inactivator, C3B/C4B inactivator, Complement component I, Complement component factor i, Factor I, FI, KAF, I factor, I factor (complement), C3b-INA, Complement factor I, Light chain of factor I

CFI

Target/Specificity Human 65 kD protein

**Reconstitution & Storage** 

**Antiserum** 

### **Precautions**

Anti-CFI / Complement Factor I Antibody (Internal) is for research use only and not for use in diagnostic or therapeutic procedures.

# Anti-CFI / Complement Factor I Antibody (Internal) - Protein Information

**Name CFI** 

Synonyms IF

## **Function**

Trypsin-like serine protease that plays an essential role in regulating the immune response by controlling all complement pathways. Inhibits these pathways by cleaving three peptide bonds in the alpha- chain of C3b and two bonds in the alpha-chain of C4b thereby inactivating these proteins (PubMed:<a href="http://www.uniprot.org/citations/17320177" target="\_blank">17320177</a>, PubMed:<a href="http://www.uniprot.org/citations/7360115"

target="\_blank">7360115</a>). Essential cofactors for these reactions include factor H and C4BP



in the fluid phase and membrane cofactor protein/CD46 and CR1 on cell surfaces (PubMed:<a href="http://www.uniprot.org/citations/12055245" target="\_blank">12055245</a>, PubMed:<a href="http://www.uniprot.org/citations/2141838" target="\_blank">2141838</a>, PubMed:<a href="http://www.uniprot.org/citations/9605165" target="\_blank">9605165</a>). The presence of these cofactors on healthy cells allows degradation of deposited C3b by CFI in order to prevent undesired complement activation, while in apoptotic cells or microbes, the absence of such cofactors leads to C3b-mediated complement activation and subsequent opsonization (PubMed:<a href="http://www.uniprot.org/citations/28671664" target=" blank">28671664</a>).

## **Cellular Location**

Secreted, extracellular space. Secreted

#### **Tissue Location**

Expressed in the liver by hepatocytes (PubMed:6327681). Also present in other cells such as monocytes, fibroblasts or keratinocytes (PubMed:17320177, PubMed:6444659)

# Anti-CFI / Complement Factor I Antibody (Internal) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-CFI / Complement Factor I Antibody (Internal) - Images