

Anti-TLR9 Antibody (clone 72-1665)
Rat Anti Human Monoclonal Antibody
Catalog # ALS17763

Specification

Anti-TLR9 Antibody (clone 72-1665) - Product Information

Application	WB, IHC-P, FC
Primary Accession	Q9NR96
Predicted	Human
Host	Rat
Clonality	Monoclonal
Isotype	IgG2a,k
Calculated MW	115860
Dilution	WB~~1:1000 IHC-P~~N/A FC~~1:10~50

Anti-TLR9 Antibody (clone 72-1665) - Additional Information

Gene ID 54106

Alias Symbol TLR9

Other Names

TLR9, CD289, CD289 antigen, Scri2a, Toll-like receptor 9

Reconstitution & Storage

Affinity purified

Precautions

Anti-TLR9 Antibody (clone 72-1665) is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-TLR9 Antibody (clone 72-1665) - Protein Information

Name TLR9

Function

Key component of innate and adaptive immunity. TLRs (Toll-like receptors) control host immune response against pathogens through recognition of molecular patterns specific to microorganisms. TLR9 is a nucleotide-sensing TLR which is activated by unmethylated cytidine-phosphate-guanosine (CpG) dinucleotides (PubMed:14716310). Acts via MYD88 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response (PubMed:11564765, PubMed:17932028). Controls lymphocyte response to Helicobacter infection (By similarity). Upon CpG stimulation, induces B-cell proliferation, activation, survival and antibody production (PubMed:23857366)

target="_blank">23857366).

Cellular Location

Endoplasmic reticulum membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q9EQU3}. Early endosome membrane. Lysosome {ECO:0000250|UniProtKB:Q9EQU3} Cytoplasmic vesicle, phagosome {ECO:0000250|UniProtKB:Q9EQU3}. Golgi apparatus membrane. Note=Relocalizes from endoplasmic reticulum to endosome and lysosome upon stimulation with agonist. Exit from the ER requires UNC93B1. Endolysosomal localization is required for proteolytic cleavage and subsequent activation Intracellular localization of the active receptor may prevent from responding to self nucleic acid. {ECO:0000250|UniProtKB:Q9EQU3, ECO:0000269|PubMed:14716310, ECO:0000269|PubMed:38169466}

Tissue Location

Highly expressed in spleen, lymph node, tonsil and peripheral blood leukocytes, especially in plasmacytoid pre-dendritic cells. Levels are much lower in monocytes and CD11c+ immature dendritic cells. Also detected in lung and liver

Anti-TLR9 Antibody (clone 72-1665) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TLR9 Antibody (clone 72-1665) - Images