

Anti-TRIM11 Antibody (aa249-263)
Goat Anti Human Polyclonal Antibody
Catalog # ALS17872**Specification**

Anti-TRIM11 Antibody (aa249-263) - Product Information

Application	WB, IHC-P, E
Primary Accession	Q96F44
Predicted	Human, Monkey
Host	Goat
Clonality	Polyclonal
Calculated MW	52774

Anti-TRIM11 Antibody (aa249-263) - Additional Information**Gene ID** 81559**Alias Symbol** **TRIM11**
Other Names

TRIM11, BIA1, Protein BIA1, RNF92, RING finger protein 92, Tripartite motif-containing 11, Tripartite motif containing 11

Target/Specificity
Human TRIM11.**Reconstitution & Storage**
Immunoaffinity purified**Precautions**
Anti-TRIM11 Antibody (aa249-263) is for research use only and not for use in diagnostic or therapeutic procedures.**Anti-TRIM11 Antibody (aa249-263) - Protein Information****Name** TRIM11 {ECO:0000303|PubMed:16904669, ECO:0000312|HGNC:HGNC:16281}**Function**

E3 ubiquitin-protein ligase that promotes the degradation of insoluble ubiquitinated proteins, including insoluble PAX6, poly-Gln repeat expanded HTT and poly-Ala repeat expanded ARX (By similarity). Mediates PAX6 ubiquitination leading to proteasomal degradation, thereby modulating cortical neurogenesis (By similarity). May also inhibit PAX6 transcriptional activity, possibly in part by preventing the binding of PAX6 to its consensus sequences (By similarity). May contribute to the regulation of the intracellular level of HN (humanin) or HN-containing proteins through the proteasomal degradation pathway (By similarity). Mediates MED15 ubiquitination leading to proteasomal degradation (PubMed:16904669). May contribute to the innate restriction of retroviruses (PubMed:18248090). Upon overexpression, reduces HIV-1 and murine leukemia virus infectivity, by suppressing viral

gene expression (PubMed:18248090). Antiviral activity depends on a functional E3 ubiquitin-protein ligase domain (PubMed:18248090). May regulate TRIM5 turnover via the proteasome pathway, thus counteracting the TRIM5-mediated cross-species restriction of retroviral infection at early stages of the retroviral life cycle (PubMed:18248090). Acts as an inhibitor of the AIM2 inflammasome by promoting autophagy-dependent degradation of AIM2 (PubMed:27498865). Mechanistically, undergoes autoubiquitination upon DNA stimulation, promoting interaction with AIM2 and SQSTM1/p62, leading to AIM2 recruitment to autophagosomes (PubMed:27498865).

Cellular Location

Cytoplasm. Nucleus

Tissue Location

Ubiquitous..

Anti-TRIM11 Antibody (aa249-263) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TRIM11 Antibody (aa249-263) - Images