

### **Anti-WTAP Antibody (clone 1B11)**

Mouse Anti Human Monoclonal Antibody Catalog # ALS17991

### **Specification**

## Anti-WTAP Antibody (clone 1B11) - Product Information

Application WB, IHC-P, E
Primary Accession Q15007
Predicted Human
Host Mouse
Clonality Monoclonal
Isotype IgG2a,k
Calculated MW 44244

# Anti-WTAP Antibody (clone 1B11) - Additional Information

**Gene ID 9589** 

Alias Symbol WTAP

**Other Names** 

WTAP, HFL(2)D, KIAA0105, PNAS-132, WT1-associated protein, Female-lethal(2)D homolog

**Target/Specificity** 

**Human WTAP** 

**Reconstitution & Storage** 

Protein A purified

#### **Precautions**

Anti-WTAP Antibody (clone 1B11) is for research use only and not for use in diagnostic or therapeutic procedures.

### Anti-WTAP Antibody (clone 1B11) - Protein Information

Name WTAP {ECO:0000303|PubMed:11001926, ECO:0000312|HGNC:HGNC:16846}

### **Function**

Associated component of the WMM complex, a complex that mediates N6-methyladenosine (m6A) methylation of RNAs, a modification that plays a role in the efficiency of mRNA splicing and RNA processing (PubMed:<a href="http://www.uniprot.org/citations/29507755" target="blank">29507755</a>). Required for accumulation of METTL13 and METTL14 to nuclear

speckle (PubMed:<a href="http://www.uniprot.org/citations/24316715"

target="\_blank">24316715</a>, PubMed:<a href="http://www.uniprot.org/citations/24407421" target=" blank">24407421</a>, PubMed:<a href="http://www.uniprot.org/citations/24981863"

target="blank">24981863</a>). Acts as a mRNA splicing regulator (PubMed:<a

href="http://www.uniprot.org/citations/12444081" target="\_blank">12444081</a>). Regulates G2/M cell-cycle transition by binding to the 3' UTR of CCNA2, which enhances its stability

(PubMed:<a href="http://www.uniprot.org/citations/17088532" target=" blank">17088532</a>).



Impairs WT1 DNA-binding ability and inhibits expression of WT1 target genes (PubMed:<a href="http://www.uniprot.org/citations/17095724" target="blank">17095724</a>).

#### **Cellular Location**

Nucleus speckle. Nucleus, nucleoplasm. Cytoplasm {ECO:0000250|UniProtKB:Q9ER69}. Note=Mainly nuclear with some fraction located in the cytoplasm. ZC3H13 is required to anchor component of the MACOM subcomplex, such as VIRMA, in the nucleus {ECO:0000250|UniProtKB:Q9ER69}

**Tissue Location** 

Ubiquitously expressed.

# Anti-WTAP Antibody (clone 1B11) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Anti-WTAP Antibody (clone 1B11) - Images