

Anti-CFTR Antibody
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS18178**Specification**

Anti-CFTR Antibody - Product Information

Application	WB, IHC-P, IF, ICC, IP
Primary Accession	P13569
Predicted	Human, Guinea Pig
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG

Anti-CFTR Antibody - Additional Information**Gene ID** 1080

Alias Symbol	CFTR
Other Names	
CFTR, ABC35, ABCC7, CF, DJ760C5.1, MRP7, CFTR/MRP, TNR-CFTR	

Target/Specificity

The antibody detects cystic fibrosis transmembrane conductance factor (CFTR) from cells overexpressing the human protein. It has been successfully used in immunocytochemistry procedures. Immunocytochemical staining of HEK293 cells overexpressing huma ...

Reconstitution & Storage

Immunoaffinity purified

Precautions

Anti-CFTR Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-CFTR Antibody - Protein Information**Name** CFTR**Synonyms** ABCC7**Function**

Epithelial ion channel that plays an important role in the regulation of epithelial ion and water transport and fluid homeostasis (PubMed: [26823428](http://www.uniprot.org/citations/26823428)). Mediates the transport of chloride ions across the cell membrane (PubMed: [10792060](http://www.uniprot.org/citations/10792060), PubMed: [11524016](http://www.uniprot.org/citations/11524016), PubMed: [11707463](http://www.uniprot.org/citations/11707463), PubMed: [12519745](http://www.uniprot.org/citations/12519745), PubMed: [15010471](http://www.uniprot.org/citations/15010471))

target="_blank">15010471, PubMed:12588899, PubMed:17036051, PubMed:19398555, PubMed:19621064, PubMed:22178883, PubMed:25330774, PubMed:1712898, PubMed:8910473, PubMed:9804160, PubMed:12529365, PubMed:17182731, PubMed:26846474, PubMed:28087700). Channel activity is coupled to ATP hydrolysis (PubMed:8910473). The ion channel is also permeable to HCO₃(-); selectivity depends on the extracellular chloride concentration (PubMed:15010471, PubMed:19019741). Exerts its function also by modulating the activity of other ion channels and transporters (PubMed:12403779, PubMed:22178883, PubMed:22121115, PubMed:27941075). Plays an important role in airway fluid homeostasis (PubMed:16645176, PubMed:19621064, PubMed:26823428). Contributes to the regulation of the pH and the ion content of the airway surface fluid layer and thereby plays an important role in defense against pathogens (PubMed:14668433, PubMed:16645176, PubMed:26823428). Modulates the activity of the epithelial sodium channel (ENaC) complex, in part by regulating the cell surface expression of the ENaC complex (PubMed:17434346, PubMed:27941075, PubMed:17182731). Inhibits the activity of the ENaC channel containing subunits SCNN1A, SCNN1B and SCNN1G (PubMed:17182731). Inhibits the activity of the ENaC channel containing subunits SCNN1D, SCNN1B and SCNN1G, but not of the ENaC channel containing subunits SCNN1A, SCNN1B and SCNN1G (PubMed:17182731, PubMed:27941075). May regulate bicarbonate secretion and salvage in epithelial cells by regulating the transporter SLC4A7 (PubMed:12403779). Can inhibit the chloride channel activity of ANO1 (PubMed:22178883). Plays a role in the chloride and bicarbonate homeostasis during sperm epididymal maturation and capacitation (PubMed:19923167, PubMed:27714810).

Cellular Location

Apical cell membrane; Multi-pass membrane protein {ECO:0000269|Ref.55}. Early endosome membrane; Multi-pass membrane protein {ECO:0000269|Ref.55}. Cell membrane; Multi-pass membrane protein {ECO:0000269|Ref.55}. Recycling endosome membrane; Multi-pass membrane protein {ECO:0000269|Ref.55}. Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000269|Ref.55}. Nucleus {ECO:0000250|UniProtKB:P34158}. Note=The channel is

internalized from the cell surface into an endosomal recycling compartment, from where it is recycled to the cell membrane (PubMed:17462998, PubMed:19398555, PubMed:20008117). In the oviduct and bronchus, detected on the apical side of epithelial cells, but not associated with cilia (PubMed:22207244). In Sertoli cells, a processed product is detected in the nucleus (By similarity). ER stress induces GORASP2-mediated unconventional (ER/Golgi-independent) trafficking of core-glycosylated CFTR to cell membrane (PubMed:21884936). {ECO:0000250|UniProtKB:P34158, ECO:0000269|PubMed:19398555, ECO:0000269|PubMed:20008117, ECO:0000269|PubMed:21884936, ECO:0000269|PubMed:22207244, ECO:0000305|PubMed:17462998}

Tissue Location

Expressed in the respiratory airway, including bronchial epithelium, and in the female reproductive tract, including oviduct (at protein level) (PubMed:22207244, PubMed:15716351). Detected in pancreatic intercalated ducts in the exocrine tissue, on epithelial cells in intralobular striated ducts in sublingual salivary glands, on apical membranes of crypt cells throughout the small and large intestine, and on the reabsorptive duct in eccrine sweat glands (PubMed:1284548, PubMed:28130590). Detected on the equatorial segment of the sperm head (at protein level) (PubMed:19923167). Detected in nasal and bronchial superficial epithelium (PubMed:15716351). Expressed by the central cells on the sebaceous glands, dermal adipocytes and, at lower levels, by epithelial cells (PubMed:28130590)

Volume

Array

Anti-CFTR Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CFTR Antibody - Images