

Anti-FN1 / Fibronectin Antibody
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS18266**Specification**

Anti-FN1 / Fibronectin Antibody - Product Information

Application	WB, IHC-P, E
Primary Accession	P02751
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	272320

Anti-FN1 / Fibronectin Antibody - Additional Information**Gene ID 2335**Alias Symbol **FN1****Other Names**

FN1, CIG, Cold-insoluble globulin, Fibronectin 1, FINC, LETS, GFND, ED-B, Fibronectin, FN, FNZ, GFND2, Migration-stimulating factor, MSF

Reconstitution & Storage

Affinity purified

Precautions

Anti-FN1 / Fibronectin Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-FN1 / Fibronectin Antibody - Protein InformationName FN1 ([HGNC:3778](#))**Synonyms FN****Function**

Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin (PubMed: [3024962](http://www.uniprot.org/citations/3024962), PubMed: [3900070](http://www.uniprot.org/citations/3900070), PubMed: [3593230](http://www.uniprot.org/citations/3593230), PubMed: [7989369](http://www.uniprot.org/citations/7989369)). Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape (PubMed: [3024962](http://www.uniprot.org/citations/3024962), PubMed: [3900070](http://www.uniprot.org/citations/3900070), PubMed: [3593230](http://www.uniprot.org/citations/3593230), PubMed: [7989369](http://www.uniprot.org/citations/7989369)). Involved in

osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization (By similarity). Participates in the regulation of type I collagen deposition by osteoblasts (By similarity). Acts as a ligand for the LILRB4 receptor, inhibiting FCGR1A/CD64-mediated monocyte activation (PubMed:34089617).

Cellular Location

Secreted, extracellular space, extracellular matrix. Secreted {ECO:0000250|UniProtKB:P11276}

Tissue Location

Expressed in the inner limiting membrane and around blood vessels in the retina (at protein level) (PubMed:29777959) Plasma FN (soluble dimeric form) is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. Ugl-Y1, Ugl-Y2 and Ugl-Y3 are found in urine (PubMed:17614963).

Anti-FN1 / Fibronectin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-FN1 / Fibronectin Antibody - Images