

Anti-HTR2B / 5-HT2B Receptor Antibody
Rabbit Anti Human Polyclonal Antibody
Catalog # ALS18418

Specification

Anti-HTR2B / 5-HT2B Receptor Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P41595
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54298

Anti-HTR2B / 5-HT2B Receptor Antibody - Additional Information

Gene ID 3357

Alias Symbol HTR2B

Other Names

HTR2B, 5-HT 2B receptor, 5HT2B Receptor, 5-HT(2B), 5-HT-2B, 5-HT2b receptor, 5-HT2B, Serotonin 2b receptor, Serotonin 5-HT-2b receptor, Serotonin receptor 2B

Target/Specificity

Human HTR1B / 5-HT2B Receptor

Reconstitution & Storage

Immunoaffinity purified

Precautions

Anti-HTR2B / 5-HT2B Receptor Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Anti-HTR2B / 5-HT2B Receptor Antibody - Protein Information

Name HTR2B ([HGNC:5294](#))

Function

G-protein coupled receptor for 5-hydroxytryptamine (serotonin) (PubMed:18703043, PubMed:23519210, PubMed:7926008, PubMed:8078486, PubMed:8143856, PubMed:8882600). Also functions as a receptor for various ergot alkaloid derivatives and psychoactive substances (PubMed:12970106, PubMed:18703043, PubMed:23519210, PubMed:>23519215, PubMed:>24357322, PubMed:>28129538, PubMed:>30127358, PubMed:>36087581, PubMed:>7926008, PubMed:>8078486, PubMed:>8143856). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors (PubMed:>23519215, PubMed:>28129538, PubMed:>8078486, PubMed:>8143856, PubMed:>8882600). HTR2B is coupled to G(q)/G(11) G alpha proteins and activates phospholipase C-beta, releasing diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP3) second messengers that modulate the activity of phosphatidylinositol 3- kinase and promote the release of Ca(2+) ions from intracellular stores, respectively (PubMed:>18703043, PubMed:>23519215, PubMed:>28129538, PubMed:>30127358, PubMed:>36087581, PubMed:>8078486, PubMed:>8143856, PubMed:>8882600). Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways (PubMed:>23519215, PubMed:>28129538, PubMed:>30127358, PubMed:>36087581). Plays a role in the regulation of dopamine and 5-hydroxytryptamine release, 5-hydroxytryptamine uptake and in the regulation of extracellular dopamine and 5-hydroxytryptamine levels, and thereby affects neural activity. May play a role in the perception of pain (By similarity). Plays a role in the regulation of behavior, including impulsive behavior (PubMed:>21179162). Required for normal proliferation of embryonic cardiac myocytes and normal heart development (By similarity). Protects cardiomyocytes against apoptosis (By similarity). Plays a role in the adaptation of pulmonary arteries to chronic hypoxia (By similarity). Plays a role in vasoconstriction (By similarity). Required for normal osteoblast function and proliferation, and for maintaining normal bone density (By similarity). Required for normal proliferation of the interstitial cells of Cajal in the intestine (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Synapse, synaptosome {ECO:0000250|UniProtKB:Q02152}

Tissue Location

Ubiquitous. Detected in liver, kidney, heart, pulmonary artery, and intestine. Detected at lower levels in blood, placenta and brain, especially in cerebellum, occipital cortex and frontal cortex.

Anti-HTR2B / 5-HT2B Receptor Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-HTR2B / 5-HT2B Receptor Antibody - Images