

**Anti-DDX39B / UAP56 Antibody (aa2-251)**  
**Rabbit Anti Human Polyclonal Antibody**  
**Catalog # ALS18515****Specification**

---

**Anti-DDX39B / UAP56 Antibody (aa2-251) - Product Information**

Application	WB, IHC-P, E
Primary Accession	<a href="#">Q13838</a>
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	48991

**Anti-DDX39B / UAP56 Antibody (aa2-251) - Additional Information****Gene ID** 7919Alias Symbol **DDX39B****Other Names**

DDX39B, BAT1, DEAD box protein UAP56, HLA-B associated transcript 1, UAP56, ATP-dependent RNA helicase p47, D6S81E, Spliceosome RNA helicase BAT1

**Target/Specificity**

Human DDX39B / UAP56 / BAT1

**Reconstitution & Storage**

Caprylic acid and ammonium sulfate precipitation

**Precautions**

Anti-DDX39B / UAP56 Antibody (aa2-251) is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-DDX39B / UAP56 Antibody (aa2-251) - Protein Information****Name** DDX39B ([HGNC:13917](#))**Synonyms** BAT1, UAP56**Function**

Involved in nuclear export of spliced and unspliced mRNA. Assembling component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced pre-mRNA. TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and cap-dependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP/NFX1 pathway. May undergo several rounds of ATP hydrolysis during assembly of TREX to drive subsequent loading of components such as ALYREF/THOC and CHTOP onto mRNA. Also associates

with pre-mRNA independent of ALYREF/THOC4 and the THO complex. Involved in the nuclear export of intronless mRNA; the ATP-bound form is proposed to recruit export adapter ALYREF/THOC4 to intronless mRNA; its ATPase activity is cooperatively stimulated by RNA and ALYREF/THOC4 and ATP hydrolysis is thought to trigger the dissociation from RNA to allow the association of ALYREF/THOC4 and the NXF1-NXT1 heterodimer. Involved in transcription elongation and genome stability.

**Cellular Location**

Nucleus. Nucleus speckle. Cytoplasm. Note=Can translocate to the cytoplasm in the presence of MX1. TREX complex assembly seems to occur in regions surrounding nuclear speckles known as perispeckles

**Anti-DDX39B / UAP56 Antibody (aa2-251) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-DDX39B / UAP56 Antibody (aa2-251) - Images**