

**Anti-PPP1CA / PP1-Alpha Antibody**  
**Rabbit Anti Human Polyclonal Antibody**  
**Catalog # ALS18590****Specification**

---

**Anti-PPP1CA / PP1-Alpha Antibody - Product Information**

Application	WB, IHC-P, IF
Primary Accession	<a href="#">P62136</a>
Predicted	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Calculated MW	37512

**Anti-PPP1CA / PP1-Alpha Antibody - Additional Information****Gene ID** 5499**Alias Symbol** **PPP1CA****Other Names**

PPP1CA, PP-1A, pp1C alpha, PP1 alpha, PP1-Alpha, PP1alpha, PPP1A, Protein phosphatase 1 alpha

**Target/Specificity**

Human PPP1CA / PP1-Alpha

**Reconstitution & Storage**

Affinity purified

**Precautions**

Anti-PPP1CA / PP1-Alpha Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-PPP1CA / PP1-Alpha Antibody - Protein Information****Name** PPP1CA**Synonyms** PPP1A**Function**

Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets. Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density-associated Ca(2+)/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase. Regulates NEK2 function in terms of kinase activity and centrosome number and splitting, both in the presence and absence

of radiation- induced DNA damage. Regulator of neural tube and optic fissure closure, and enteric neural crest cell (ENCCs) migration during development. In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. May dephosphorylate CSNK1D and CSNK1E. Dephosphorylates the 'Ser-418' residue of FOXP3 in regulatory T-cells (Treg) from patients with rheumatoid arthritis, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed:<a href="http://www.uniprot.org/citations/23396208" target="\_blank">23396208</a>). Dephosphorylates CENPA (PubMed:<a href="http://www.uniprot.org/citations/25556658" target="\_blank">25556658</a>). Dephosphorylates the 'Ser-139' residue of ATG16L1 causing dissociation of ATG12-ATG5-ATG16L1 complex, thereby inhibiting autophagy (PubMed:<a href="http://www.uniprot.org/citations/26083323" target="\_blank">26083323</a>).

#### **Cellular Location**

Cytoplasm. Nucleus, nucleoplasm. Nucleus, nucleolus Note=Primarily nuclear and largely excluded from the nucleolus. Highly mobile in cells and can be relocalized through interaction with targeting subunits. NOM1 plays a role in targeting this protein to the nucleolus. In the presence of PPP1R8 relocalizes from the nucleus to nuclear speckles. Shuttles toward the cytosol during infection with VEEV (PubMed:29769351).

#### **Anti-PPP1CA / PP1-Alpha Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-PPP1CA / PP1-Alpha Antibody - Images**