

#### **GST Antibody**

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM1011b

## **Specification**

#### **GST Antibody - Product Information**

Application WB,E
Host Mouse
Clonality Monoclonal
Isotype IgG1 k
Calculated MW 26000 Da

## **GST Antibody - Additional Information**

#### **Other Names**

Glutathione S-transferase

#### **Target/Specificity**

Purified recombinant GST fusion protein was used to produced this monoclonal antibody.

#### **Dilution**

WB~~1:100~500

E~~Use at an assay dependent concentration.

#### **Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

## Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

GST Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **GST Antibody - Protein Information**

## **GST Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence



- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **GST Antibody - Images**



Western blot analysis of anti-GST Mab in recombinant GST protein. GST(arrow) was detected using the purified Mab .

## **GST Antibody - Background**

Glutathione S-transferase (GST) was originally cloned from parasite Schistosoma japonicum and it is now a widely used protein fusion partner. Vectors containing GST Tag have been developed for both prokaryotic and eukaryotic systems. The GST fusion proteins are easily purified from cell lysates by affinity chromatography using Glutathione Sepharose 4B to elute out the GST fusion protein from the column with a denaturing form of glutathione. Using the Abgent anti-GST antibody provides a simple solution to detect the expression of GST fusion proteins in cells.

# **GST Antibody - References**

Smith, D.B. and Johnson, K.S., (1988). Gene 67, 31. Parker, M.W. et al., (1990) J. Mol. Biol. 213, 221. Toye, B. et al., (1990) Infect. Immun. 58, 3909. Guan, K.L. and Dixon, J.E. (1991) Anal. Biochem. 192, 262

# **GST Antibody - Citations**

• HnRNP F/H associate with hTERC and telomerase holoenzyme to modulate telomerase function and promote cell proliferation