

HSPA1A Antibody

Mouse Monoclonal Antibody (Mab)
Catalog # AM1877b

Specification

HSPA1A Antibody - Product Information

Application IF, WB,E Primary Accession P08107

Other Accession <u>Q6S4N2</u>, <u>Q27965</u>, <u>P34930</u>, <u>Q27975</u>,

NP_005336.3, P0DMV8, P0DMV9

HSPA1A Antibody - Additional Information

Other Names

Heat shock 70 kDa protein 1A/1B, Heat shock 70 kDa protein 1/2, HSP70-1/HSP70-2, HSP701/HSP702, HSPA1A, HSPA1, HSX70

Target/Specificity

This HSPA1A antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 574-600 amino acids from human HSPA1A.

Dilution

IF~~1:10~50 WB~~1:100~2000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HSPA1A Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

HSPA1A Antibody - Protein Information


HSPA1A Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

HSPA1A Antibody - Images



Confocal immunofluorescent analysis of HSPA1A Antibody (Cat#AM1877b) with Hela cell followed by Alexa Fluor® 488-conjugated goat anti-mouse IgG (green). DAPI was used to stain the cell nuclear (blue).



HSPA1A antibody (Cat. #AM1877b) western blot analysis in HL-60 cell line lysates $(35\mu g/lane)$. This demonstrates the HSPA1A antibody detected the HSPA1A protein (arrow).

HSPA1A Antibody - Background

This intronless gene encodes a 70kDa heat shock protein which is a member of the heat shock protein 70 family. In conjuction with other heat shock proteins, this protein stabilizes





existing proteins against aggregation and mediates the folding of newly translated proteins in the cytosol and in organelles. It is also involved in the ubiquitin-proteasome pathway through interaction with the AU-rich element RNA-binding protein 1. The gene is located in the major histocompatibility complex class III region, in a cluster with two closely related genes which encode similar proteins.

HSPA1A Antibody - References

Wang, Y., et al. Clin. Chem. Lab. Med. 48(11):1657-1663(2010) Eisenberg, D.P., et al. Surgery 148(2):325-334(2010) Rusai, K., et al. Transplant. Proc. 42(6):2309-2311(2010) Ebrahimi, M., et al. Mol. Vis. 16, 1680-1688 (2010) : Lanneau, D., et al. ScientificWorldJournal 10, 1543-1552 (2010) :