

YEATS4 antibody (Ascites)

Mouse Monoclonal Antibody (Mab)
Catalog # AM1885a

Specification

YEATS4 antibody (Ascites) - Product Information

Application WB,E
Primary Accession O95619
Reactivity Human
Host Mouse
Clonality Monoclonal
Isotype IgG1,K

YEATS4 antibody (Ascites) - Additional Information

Gene ID 8089

Other Names

YEATS domain-containing protein 4, Glioma-amplified sequence 41, Gas41, NuMA-binding protein 1, NuBI-1, NuBI1, YEATS4, GAS41

Target/Specificity

This YEATS4 monoclonal antibody is generated from mouse immunized with YEATS4 recombinant protein.

Dilution

WB~~1:100~16000

Format

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

YEATS4 antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

YEATS4 antibody (Ascites) - Protein Information

Name YEATS4 (HGNC:24859)

Function Chromatin reader component of the NuA4 histone acetyltransferase (HAT) complex, a complex involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A (PubMed:12963728, PubMed:14966270). Specifically recognizes and binds acylated histone H3, with a preference for histone H3 diacetylated at 'Lys-18' and 'Lys-27' (H3K18ac and H3K27ac) or histone H3 diacetylated at 'Lys-14' and 'Lys-27' (H3K14ac and



H3K27ac) (PubMed: 29437725, PubMed: 30071723, PubMed: 29900004). Also able to recognize and bind crotonylated histone H3 (PubMed: 30071723). May also recognize and bind histone H3 succinylated at 'Lys-122' (H3K122succ); additional evidences are however required to confirm this result in vivo (PubMed: 29463709). Plays a key role in histone variant H2AZ1/H2A.Z deposition into specific chromatin regions: recognizes and binds H3K14ac and H3K27ac on the promoters of actively transcribed genes and recruits NuA4-related complex to deposit H2AZ1/H2A.Z (PubMed: 29437725). H2AZ1/H2A.Z deposition is required for maintenance of embryonic stem cell (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00376, ECO:0000269|PubMed:10913114, ECO:0000269|PubMed:18445686}

Tissue Location


Expressed in brain, heart, kidney, liver, lung, pancreas, placenta and skeletal muscle.

YEATS4 antibody (Ascites) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

YEATS4 antibody (Ascites) - Images



YEATS4 antibody (Cat. #AM1885a) western blot analysis in 293 cell line lysates ($35\mu g$ /lane). This demonstrates the YEATS4 antibody detected the YEATS4 protein (arrow).

YEATS4 antibody (Ascites) - Background

The protein encoded by this gene is found in the nucleoli. It has high sequence homology to human MLLT1, and yeast and human MLLT3 proteins. Both MLLT1 and MLLT3 proteins belong to a class of transcription factors, indicating that the encoded protein might also represent a transcription factor. This protein is thought to be required for RNA transcription. This gene has been shown to be



Tel: 858.875.1900 Fax: 858.875.1999



amplified in tumors.

YEATS4 antibody (Ascites) - References

Heisel, S., et al. BMC Mol. Biol. 11, 53 (2010): Turner, S.T., et al. Hypertension 52(2):359-365(2008) Italiano, A., et al. Int. J. Cancer 122(10):2233-2241(2008) Gudbjartsson, D.F., et al. Nat. Genet. 40(5):609-615(2008) Park, J.H., et al. Mol. Cell. Biol. 26(11):4006-4016(2006)