

**ATG4D Antibody**  
**Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM1896B****Specification**

---

**ATG4D Antibody - Product Information**

Application	IF, WB,E
Primary Accession	<a href="#">Q86TL0</a>
Other Accession	<a href="#">NP_116274.3</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1,K

**ATG4D Antibody - Additional Information****Gene ID** 84971**Other Names**

Cysteine protease ATG4D, 3422-, AUT-like 4 cysteine endopeptidase, Autophagin-4, Autophagy-related cysteine endopeptidase 4, Autophagy-related protein 4 homolog D, Cysteine protease ATG4D, mitochondrial, ATG4D, APG4D, AUTL4

**Target/Specificity**

This ATG4D monoclonal antibody is generated from mouse immunized with ATG4D recombinant protein.

**Dilution**

IF~~1:25

WB~~1:4000

E~~Use at an assay dependent concentration.

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ATG4D Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**ATG4D Antibody - Protein Information****Name** ATG4D {ECO:0000303|PubMed:19549685, ECO:0000312|HGNC:HGNC:20789}**Function** [Cysteine protease ATG4D]: Cysteine protease that plays a key role in autophagy by

mediating both proteolytic activation and delipidation of ATG8 family proteins (PubMed:[21177865](#), PubMed:[29458288](#), PubMed:[30661429](#)). The protease activity is required for proteolytic activation of ATG8 family proteins: cleaves the C-terminal amino acid of ATG8 proteins MAP1LC3 and GABARAPL2, to reveal a C-terminal glycine (PubMed:[21177865](#)). Exposure of the glycine at the C-terminus is essential for ATG8 proteins conjugation to phosphatidylethanolamine (PE) and insertion to membranes, which is necessary for autophagy (By similarity). In addition to the protease activity, also mediates delipidation of ATG8 family proteins (PubMed:[29458288](#), PubMed:[33909989](#)). Catalyzes delipidation of PE-conjugated forms of ATG8 proteins during macroautophagy (PubMed:[29458288](#), PubMed:[33909989](#)). Also involved in non-canonical autophagy, a parallel pathway involving conjugation of ATG8 proteins to single membranes at endolysosomal compartments, by catalyzing delipidation of ATG8 proteins conjugated to phosphatidylserine (PS) (PubMed:[33909989](#)). ATG4D plays a role in the autophagy-mediated neuronal homeostasis in the central nervous system (By similarity). Compared to other members of the family (ATG4A, ATG4B or ATG4C), constitutes the major protein for the delipidation activity, while it promotes weak proteolytic activation of ATG8 proteins (By similarity). Involved in phagophore growth during mitophagy independently of its protease activity and of ATG8 proteins: acts by regulating ATG9A trafficking to mitochondria and promoting phagophore- endoplasmic reticulum contacts during the lipid transfer phase of mitophagy (PubMed:[33773106](#)).

#### Cellular Location

[Cysteine protease ATG4D]: Cytoplasm

#### Tissue Location


Widely expressed in testis.

### ATG4D Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### ATG4D Antibody - Images



Immunofluorescent analysis of U251 cells, using ATG4D Antibody(Cat. #AM1896b). U251

cells(right) were treated with Chloroquine (50  $\mu$ M,16h). AM1896b was diluted at 1:25 dilution. Dylight Fluor 488-conjugated goat anti-mouse IgG at 1:400 dilution was used as the secondary antibody (green).DAPI was used to stain the cell nuclear (blue).



Anti-ATG4D at 1:4000 dilution + A431 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 53 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

#### **ATG4D Antibody - Background**

Autophagy is the process by which endogenous proteins and damaged organelles are destroyed intracellularly. Autophagy is postulated to be essential for cell homeostasis and cell remodeling during differentiation, metamorphosis, non-apoptotic cell death, and aging. Reduced levels of autophagy have been described in some malignant tumors, and a role for autophagy in controlling the unregulated cell growth linked to cancer has been proposed. This gene encodes a member of the autophagin protein family. The encoded protein is also designated as a member of the C-54 family of cysteine proteases.

#### **ATG4D Antibody - References**

Betin, V.M., et al. J. Cell. Sci. 122 (PT 14), 2554-2566 (2009) :  
Lamesch, P., et al. Genomics 89(3):307-315(2007)  
Marino, G., et al. J. Biol. Chem. 278(6):3671-3678(2003)