

INS Antibody (Ascites)
Mouse Monoclonal Antibody (Mab)
Catalog # AM1985a**Specification**

INS Antibody (Ascites) - Product Information

Application	WB,E
Primary Accession	P01308
Other Accession	NP_001172027.1 , NP_000198.1 , NP_001172026.1
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgM
Calculated MW	11981
Antigen Region	35-64

INS Antibody (Ascites) - Additional Information**Gene ID** 3630**Other Names**

Insulin, Insulin B chain, Insulin A chain, INS

Target/Specificity

This INS antibody is generated from mice immunized with a KLH conjugated synthetic peptide between 35-64 amino acids from human INS.

Dilution

WB~~1:1000~8000

E~~Use at an assay dependent concentration.

Format

Mouse monoclonal antibody supplied in crude ascites with 0.09% (W/V) sodium azide.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

INS Antibody (Ascites) is for research use only and not for use in diagnostic or therapeutic procedures.

INS Antibody (Ascites) - Protein Information**Name** INS**Function** Insulin decreases blood glucose concentration. It increases cell permeability to

monosaccharides, amino acids and fatty acids. It accelerates glycolysis, the pentose phosphate cycle, and glycogen synthesis in liver.

Cellular Location


Secreted.

INS Antibody (Ascites) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

INS Antibody (Ascites) - Images



INS Antibody (Cat. #AM1985a) western blot analysis in CEM cell line lysates (35µg/lane). This demonstrates the INS antibody detected the INS protein (arrow).

INS Antibody (Ascites) - Background

After removal of the precursor signal peptide, proinsulin is post-translationally cleaved into three peptides: the B chain and A chain peptides, which are covalently linked via two disulfide bonds to form insulin, and C-peptide. Binding of insulin to the insulin receptor (INSR) stimulates glucose uptake. A multitude of mutant alleles with phenotypic effects have been identified. There is a read-through gene, INS-IGF2, which overlaps with this gene at the 5' region and with the IGF2 gene at the 3' region. Alternative splicing results in multiple transcript variants. [provided by RefSeq].

INS Antibody (Ascites) - References

Hinks, A., et al. Ann. Rheum. Dis. 69(12):2169-2172(2010)

Breuer, T.G., et al. Eur. J. Endocrinol. 163(4):551-558(2010)
Andersen, M.K., et al. Diabetes Care 33(9):2062-2064(2010)
Ferron, M., et al. Cell 142(2):296-308(2010)
Authier, F., et al. J. Biol. Chem. 277(11):9437-9446(2002)