

UCH37 (UCHL5) Antibody
Mouse Monoclonal Antibody (Mab)
Catalog # AM2200b**Specification**

UCH37 (UCHL5) Antibody - Product Information

Application	WB,E
Primary Accession	Q9Y5K5
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1

UCH37 (UCHL5) Antibody - Additional Information**Gene ID** 51377**Other Names**

Ubiquitin carboxyl-terminal hydrolase isozyme L5, UCH-L5, Ubiquitin C-terminal hydrolase UCH37, Ubiquitin thioesterase L5, UCHL5, UCH37

Target/Specificity

Purified His-tagged UCH37 protein was used to produced this monoclonal antibody.

Dilution

WB~~1:2000-1:4000

E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UCH37 (UCHL5) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

UCH37 (UCHL5) Antibody - Protein Information**Name** UCHL5**Synonyms** UCH37

Function Protease that specifically cleaves 'Lys-48'-linked polyubiquitin chains. Deubiquitinating enzyme associated with the 19S regulatory subunit of the 26S proteasome. Putative regulatory

component of the INO80 complex; however is inactive in the INO80 complex and is activated by a transient interaction of the INO80 complex with the proteasome via ADRM1.

Cellular Location


Cytoplasm. Nucleus. Note=Associates with the proteasome 19S subunit in the cytoplasm. Associates with the INO80 complex in the nucleus

UCH37 (UCHL5) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UCH37 (UCHL5) Antibody - Images



All lanes : Anti-UCH37 (UCHL5) at 1:2000-1:4000 dilution Lane 1: 293 whole cell lysate Lane 2: NCI-H460 whole cell lysate Lane 3: HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 37 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

UCH37 (UCHL5) Antibody - Background

Protease that specifically cleaves 'Lys-48'-linked polyubiquitin chains. Deubiquitinating enzyme associated with the 19S regulatory subunit of the 26S proteasome. Putative regulatory component of the INO80 complex; however is inactive in the INO80 complex and is activated by a transient interaction of the INO80 complex with the proteasome via ADRM1.

UCH37 (UCHL5) Antibody - References

Yao T., et al. Nat. Cell Biol. 8:994-1002(2006).

Lai C.-H., et al. Genome Res. 10:703-713(2000).

Hu R.-M., et al. Proc. Natl. Acad. Sci. U.S.A. 97:9543-9548(2000).

Kalnine N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.

Gregory S.G., et al. Nature 441:315-321(2006).