

Myoglobin Antibody (C-term)
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8438c

Specification

Myoglobin Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P02144
Other Accession	P02189 , P02150 , P68082
Reactivity	Mouse
Predicted	Horse, Monkey, Pig
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1

Myoglobin Antibody (C-term) - Additional Information

Gene ID 4151

Other Names
Myoglobin, MB

Target/Specificity

This Myoglobin antibody is generated from a mouse immunized with a KLH conjugated synthetic peptide between 118-150 amino acids from the C-terminal region of human Myoglobin.

Dilution

WB~~1:1000
E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Myoglobin Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Myoglobin Antibody (C-term) - Protein Information

Name MB ([HGNC:6915](#))

Function Monomeric heme protein which primary function is to store oxygen and facilitate its diffusion within muscle tissues. Reversibly binds oxygen through a pentacoordinated heme iron

and enables its timely and efficient release as needed during periods of heightened demand (PubMed:[30918256](#), PubMed:[34679218](#)). Depending on the oxidative conditions of tissues and cells, and in addition to its ability to bind oxygen, it also has a nitrite reductase activity whereby it regulates the production of bioactive nitric oxide (PubMed:[32891753](#)). Under stress conditions, like hypoxia and anoxia, it also protects cells against reactive oxygen species thanks to its pseudoperoxidase activity (PubMed:[34679218](#)).

Cellular Location


Cytoplasm, sarcoplasm

Myoglobin Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Myoglobin Antibody (C-term) - Images



Western blot analysis of lysates from mouse heart and skeletal muscle tissue lysates (from left to right), using Myoglobin Antibody (C-term)(Cat. #AM8438c). AM8438c was diluted at 1:1000 at each lane. A goat anti-mouse IgG H&L(HRP) at 1:3000 dilution was used as the secondary antibody. Lysates at 35µg per lane.

Myoglobin Antibody (C-term) - Background

Serves as a reserve supply of oxygen and facilitates the movement of oxygen within muscles.

Myoglobin Antibody (C-term) - References

Weller P.,et al.EMBO J. 3:439-446(1984).

Akaboshi E., et al. Gene 33:241-249(1985).
Collins J.E., et al. Genome Biol. 5:R84.1-R84.11(2004).
Ebert L., et al. Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Dunham I., et al. Nature 402:489-495(1999).