

Zap70 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8462b**Specification**

Zap70 Antibody - Product Information

Application	WB, FC, IF,E
Primary Accession	P43404
Reactivity	Human, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG2a,k
Calculated MW	70112

Zap70 Antibody - Additional Information**Gene ID** 22637**Other Names**

Tyrosine-protein kinase ZAP-70, 70 kDa zeta-chain associated protein, Syk-related tyrosine kinase, Zap70, Srk, Zap-70

Target/Specificity

This Zap70 antibody is generated from a mouse immunized with a recombinant protein.

Dilution

WB~~1:2000

FC~~1:25

IF~~1:25

E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Zap70 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Zap70 Antibody - Protein Information**Name** Zap70**Synonyms** Srk, Zap-70

Function Tyrosine kinase that plays an essential role in regulation of the adaptive immune response. Regulates motility, adhesion and cytokine expression of mature T-cells, as well as thymocyte development. Also contributes to the development and activation of primary B-lymphocytes. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylations lead to the recruitment of ZAP70 to the doubly phosphorylated TCR component CD3Z through ITAM motif at the plasma membrane. This recruitment serves to localization to the stimulated TCR and to relieve its autoinhibited conformation. Release of ZAP70 active conformation is further stabilized by phosphorylation mediated by LCK. Subsequently, ZAP70 phosphorylates at least 2 essential adapter proteins: LAT and LCP2. In turn, a large number of signaling molecules are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation. Furthermore, ZAP70 controls cytoskeleton modifications, adhesion and mobility of T- lymphocytes, thus ensuring correct delivery of effectors to the APC. ZAP70 is also required for TCR-CD3Z internalization and degradation through interaction with the E3 ubiquitin-protein ligase CBL and adapter proteins SLA and SLA2. Thus, ZAP70 regulates both T-cell activation switch on and switch off by modulating TCR expression at the T-cell surface. During thymocyte development, ZAP70 promotes survival and cell-cycle progression of developing thymocytes before positive selection (when cells are still CD4/CD8 double negative). Additionally, ZAP70-dependent signaling pathway may also contribute to primary B- cells formation and activation through B-cell receptor (BCR).

Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein. Note=In quiescent T-lymphocytes, ZAP70 is cytoplasmic. Upon TCR activation, it is recruited at the plasma membrane by interacting with CD3Z. Colocalizes together with RHOH in the immunological synapse RHOH is required for its proper localization to the cell membrane and cytoskeleton fractions in the thymocytes

Tissue Location

Isoform 1 and isoform 2 are expressed in thymus, spleen and lymph nodes.


Zap70 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Zap70 Antibody - Images






Immunofluorescent analysis of 4% paraformaldehyde-fixed, 0.1% Triton X-100 permeabilized HeLa (human cervical epithelial adenocarcinoma cell line) cells labeling Zap70 with AM8462b at 1/25 dilution, followed by Dylight® 488-conjugated goat anti-mouse IgG (NA166821) secondary antibody at 1/200 dilution (green). Immunofluorescence image showing cytoplasm staining on HeLa cell line. Cytoplasmic actin is detected with Dylight® 554 Phalloidin (PD18466410) at 1/100 dilution (red). The nuclear counter stain is DAPI (blue).



All lanes : Anti-Zap70 Antibody at 1:2000 dilution Lane 1: Jurkat whole cell lysates Lane 2: MOLT-4 whole cell lysates Lane 3: mouse thymus lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 70 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing Jurkat cells stained with AM8462b (green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AM8462b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(NA168821)) at 1/400 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG2a (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10,000 events was performed.

Zap70 Antibody - Background

Tyrosine kinase that plays an essential role in regulation of the adaptive immune response. Regulates motility, adhesion and cytokine expression of mature T-cells, as well as thymocyte development. Contributes also to the development and activation of primary B-lymphocytes. When antigen presenting cells (APC) activate T-cell receptor (TCR), a series of phosphorylations lead to the recruitment of ZAP70 to the doubly phosphorylated TCR component CD3Z through ITAM motif at the plasma membrane. This recruitment serves to localization to the stimulated TCR and to relieve its autoinhibited conformation. Release of ZAP70 active conformation is further stabilized by phosphorylation mediated by LCK. Subsequently, ZAP70 phosphorylates at least 2 essential adapter proteins: LAT and LCP2. In turn, a large number of signaling molecules are recruited and ultimately lead to lymphokine production, T-cell proliferation and differentiation. Furthermore, ZAP70 controls cytoskeleton modifications, adhesion and mobility of T-lymphocytes, thus ensuring correct delivery of effectors to the APC. ZAP70 is also required for TCR-CD3Z internalization and degradation through interaction with the E3 ubiquitin-protein ligase CBL and adapter proteins SLA and SLA2. Thus, ZAP70 regulates both T-cell activation switch on and switch off by modulating TCR expression at the T-cell surface. During thymocyte development, ZAP70 promotes survival and cell-cycle progression of developing thymocytes before positive selection (when cells are still CD4/CD8 double negative). Additionally, ZAP70-dependent signaling pathway may also contribute to primary B-cells formation and activation through B-cell receptor (BCR).

Zap70 Antibody - References

Gauen L.K.T., et al. Mol. Cell. Biol. 14:3729-3741(1994).
Wiest D.L., et al. Immunity 6:663-671(1997).
Kuroyama H., et al. Biochem. Biophys. Res. Commun. 315:935-941(2004).
Ikeda T., et al. Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.
Carninci P., et al. Science 309:1559-1563(2005).