

RAB18 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8510b**Specification**

RAB18 Antibody - Product Information

Application	WB, FC,E
Primary Accession	Q9NP72
Other Accession	Q5R5H5
Reactivity	Human, Mouse, Rat
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Antigen Region	1-206aa

RAB18 Antibody - Additional Information**Gene ID** 22931**Other Names**

Ras-related protein Rab-18, RAB18

Target/Specificity

This RAB18 antibody is generated from a mouse immunized with a recombinant protein of human RAB18.

Dilution

WB~~1:500-1:2000

FC~~1:25

E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAB18 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

RAB18 Antibody - Protein Information**Name** RAB18 ([HGNC:14244](#))**Function** The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes (PubMed:[24891604](#),

PubMed:[30970241](#)). Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (PubMed:[24891604](#), PubMed:[30970241](#)). RAB18 is required for the localization of ZFYVE1 to lipid droplets and for its function in mediating the formation of endoplasmic reticulum-lipid droplets (ER-LD) contacts (PubMed:[30970241](#)). Also required for maintaining endoplasmic reticulum structure (PubMed:[24891604](#)). Plays a role in apical endocytosis/recycling (By similarity). Plays a key role in eye and brain development and neurodegeneration (PubMed:[21473985](#)).

Cellular Location

Endoplasmic reticulum membrane. Golgi apparatus, cis-Golgi network membrane Lipid droplet. Apical cell membrane {ECO:0000250|UniProtKB:P35293}. Note=Localized to the ER membrane as well as to the cis-Golgi in fibroblasts.

Tissue Location


Ubiquitous.

RAB18 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAB18 Antibody - Images



All lanes : Anti-RAB18 Antibody at 1:500-1:2000 dilution Lane 1: PC-3 whole cell lysate Lane 2: mouse brain lysate Lane 3: rat brain lysate Lane 4: human brain lysate Lane 5: human testis lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 23 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Overlay histogram showing PC-3 cells stained with AM8510b(green line). The cells were fixed with 2% paraformaldehyde (10 min). The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AM8510b, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Mouse IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(OJ192088) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was mouse IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

RAB18 Antibody - Background

Plays a role in apical endocytosis/recycling. May be implicated in transport between the plasma membrane and early endosomes. Plays a key role in eye and brain development and neurodegeneration.

RAB18 Antibody - References

Chikri M.M.,et al.Submitted (APR-2000) to the EMBL/GenBank/DDBJ databases.
Schaefer U.,et al.FEBS Lett. 466:148-154(2000).
Dou T.,et al.DNA Seq. 16:230-234(2005).
Cui W.C.,et al.Submitted (JUL-2003) to the EMBL/GenBank/DDBJ databases.
Puhl H.L. III,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.