

## **TPM3 Antibody**

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8540b

# **Specification**

## **TPM3 Antibody - Product Information**

Application WB,E
Primary Accession P06753
Reactivity Human, Mouse
Host Mouse
Clonality monoclonal
Isotype IgG1,k
Calculated MW 32950

# **TPM3 Antibody - Additional Information**

### **Gene ID 7170**

#### **Other Names**

Tropomyosin alpha-3 chain, Gamma-tropomyosin, Tropomyosin-3, Tropomyosin-5, hTM5, TPM3

## Target/Specificity

This TPM3 antibody is generated from a mouse immunized with a recombinant protein of human TPM3.

### **Dilution**

WB~~1:2000

E~~Use at an assay dependent concentration.

#### **Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

# Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

TPM3 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

# **TPM3 Antibody - Protein Information**

### Name TPM3

**Function** Binds to actin filaments in muscle and non-muscle cells. Plays a central role, in association with the troponin complex, in the calcium dependent regulation of vertebrate striated muscle contraction. Smooth muscle contraction is regulated by interaction with caldesmon. In non-muscle cells is implicated in stabilizing cytoskeleton actin filaments.




**Cellular Location** Cytoplasm, cytoskeleton.

# **TPM3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **TPM3 Antibody - Images**



All lanes : Anti-TPM3 Antibody at 1:2000 dilution Lane 1: Hela whole cell lysate Lane 2: NIH/3T3 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 33 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

# **TPM3 Antibody - Background**

Binds to actin filaments in muscle and non-muscle cells. Plays a central role, in association with the troponin complex, in the calcium dependent regulation of vertebrate striated muscle contraction. Smooth muscle contraction is regulated by interaction with caldesmon. In non-muscle cells is implicated in stabilizing cytoskeleton actin filaments.

# **TPM3 Antibody - References**

Reinach F.C.,et al.Nature 322:648-650(1986). McLeod A.R.,et al.Nucleic Acids Res. 14:8413-8426(1986). Clayton L.,et al.J. Mol. Biol. 201:507-515(1988). Lin J.J.-C.,et al.Submitted (JUL-2000) to the EMBL/GenBank/DDBJ databases. Gregory S.G.,et al.Nature 441:315-321(2006).

