

## **RAB1B** Antibody

Purified Mouse Monoclonal Antibody (Mab) Catalog # AM8550b

### Specification

## **RAB1B Antibody - Product Information**

Application Primary Accession Other Accession Reactivity Host Clonality Isotype Calculated MW WB,E <u>O9H0U4</u> <u>O5RE13</u> Human, Mouse, Rat Mouse monoclonal IgG1,k 22171

## **RAB1B** Antibody - Additional Information

Gene ID 81876

Other Names Ras-related protein Rab-1B, RAB1B

**Target/Specificity** 

This RAB1B antibody is generated from a mouse immunized with a recombinant protein between 1-201 amino acids from human RAB1B.

**Dilution** WB~~1:4000 E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions** RAB1B Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

## **RAB1B Antibody - Protein Information**

Name RAB1B (HGNC:18370)

**Function** The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes (PubMed:<u>20545908</u>, PubMed:<u>23236136</u>). Rabs cycle between an inactive GDP-bound form and an



active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (PubMed:<u>9437002</u>). Plays a role in the initial events of the autophagic vacuole development which take place at specialized regions of the endoplasmic reticulum (PubMed:<u>20545908</u>). Regulates vesicular transport between the endoplasmic reticulum and successive Golgi compartments (By similarity). Required to modulate the compacted morphology of the Golgi (PubMed:<u>26209634</u>). Promotes the recruitment of lipid phosphatase MTMR6 to the endoplasmic reticulum- Golgi intermediate compartment (By similarity).

#### **Cellular Location**

Cytoplasm. Membrane; Lipid-anchor; Cytoplasmic side. Preautophagosomal structure membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm, perinuclear region {ECO:0000250|UniProtKB:P10536}. Note=Targeted by REP1 to membranes of specific subcellular compartments including endoplasmic reticulum, Golgi apparatus, and intermediate vesicles between these two compartments (PubMed:11389151). In the GDP-form, colocalizes with GDI in the cytoplasm (PubMed:11389151). Co-localizes with MTMR6 to the endoplasmic reticulum-Golgi intermediate compartment and to the peri- Golgi region (By similarity). {ECO:0000250|UniProtKB:P10536, ECO:0000269|PubMed:11389151}

## **RAB1B Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## RAB1B Antibody - Images



All lanes : Anti-RAB1B Antibody at 1:4000 dilution Lane 1: Hela whole cell lysate Lane 2: A431 whole cell lysate Lane 3: U-20S whole cell lysate Lane 4: rat liver lysate Lane 5: NIH/3T3 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase



conjugated at 1/10000 dilution. Predicted band size : 22 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

# RAB1B Antibody - Background

The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different set of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion. RAB1B regulates vesicular transport between the endoplasmic reticulum and successive Golgi compartments. Plays a role in the initial events of the autophagic vacuole development which take place at specialized regions of the endoplasmic reticulum.

## **RAB1B Antibody - References**

Zhao Y., et al.Submitted (SEP-1998) to the EMBL/GenBank/DDBJ databases. Wiemann S., et al.Genome Res. 11:422-435(2001). Ota T., et al.Nat. Genet. 36:40-45(2004). Bienvenut W.V., et al.Submitted (JUN-2005) to UniProtKB. Wilson A.L., et al.Biochem. J. 318:1007-1014(1996).