

PACSIN2 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8589b**Specification**

PACSIN2 Antibody - Product Information

Application	WB,E
Primary Accession	Q9UNF0
Reactivity	Human, Mouse
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Calculated MW	55739

PACSIN2 Antibody - Additional Information**Gene ID** 11252**Other Names**

Protein kinase C and casein kinase substrate in neurons protein 2, Syndapin-2, Syndapin-II, PACSIN2

Target/Specificity

This PACSIN2 antibody is generated from a mouse immunized with a recombinant protein between 250-486 amino acids from human PACSIN2.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PACSIN2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

PACSIN2 Antibody - Protein Information**Name** PACSIN2

Function Regulates the morphogenesis and endocytosis of caveolae (By similarity). Lipid-binding protein that is able to promote the tubulation of the phosphatidic acid-containing membranes it preferentially binds. Plays a role in intracellular vesicle-mediated transport. Involved in the

endocytosis of cell-surface receptors like the EGF receptor, contributing to its internalization in the absence of EGF stimulus (PubMed:[21693584](#), PubMed:[23129763](#), PubMed:[23236520](#), PubMed:[23596323](#)). Essential for endothelial organization in sprouting angiogenesis, modulates CDH5-based junctions. Facilitates endothelial front-rear polarity during migration by recruiting EHD4 and MICALL1 to asymmetric adherens junctions between leader and follower cells (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q9WVE8}. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q9WVE8}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q9WVE8}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9WVE8}; Cytoplasmic side {ECO:0000250|UniProtKB:Q9WVE8}. Cell projection, ruffle membrane {ECO:0000250|UniProtKB:Q9WVE8}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9WVE8}; Cytoplasmic side {ECO:0000250|UniProtKB:Q9WVE8}. Early endosome {ECO:0000250|UniProtKB:Q9WVE8}. Recycling endosome membrane. Cell membrane {ECO:0000250|UniProtKB:Q9WVE8}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9WVE8}; Cytoplasmic side {ECO:0000250|UniProtKB:Q9WVE8}. Cell projection. Membrane, caveola. Cell junction, adherens junction {ECO:0000250|UniProtKB:Q9WVE8}. Note=Detected at the neck of flask- shaped caveolae. Localization to tubular recycling endosomes probably requires interaction with MICALL1 and EHD1 {ECO:0000250|UniProtKB:Q9WVE8}

Tissue Location

Widely expressed.


PACSIN2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PACSIN2 Antibody - Images





All lanes : Anti-PACSIN2 Antibody at 1:2000 dilution Lane 1: Daudi whole cell lysate Lane 2: HL-60 whole cell lysate Lane 3: human brain lysate Lane 4: mouse brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 56 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

PACSIN2 Antibody - Background

Lipid-binding protein that is able to promote the tubulation of the phosphatidic acid-containing membranes it preferentially binds. Plays a role in intracellular vesicle- mediated transport. Involved in the endocytosis of cell-surface receptors like the EGF receptor, contributing to its internalization in the absence of EGF stimulus. May also play a role in the formation of caveolae at the cell membrane. Recruits DNM2 to caveolae, and thereby plays a role in caveola-mediated endocytosis.

PACSIN2 Antibody - References

Ritter B.,et al.FEBS Lett. 454:356-362(1999).
Wiemann S.,et al.Genome Res. 11:422-435(2001).
Collins J.E.,et al.Genome Biol. 5:R84.1-R84.11(2004).
Dunham I.,et al.Nature 402:489-495(1999).
Modregger J.,et al.J. Cell Sci. 113:4511-4521(2000).