

**RANBP9 Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AM8615b****Specification**

---

**RANBP9 Antibody - Product Information**

Application	WB, IHC-P,E
Primary Accession	<a href="#">Q96S59</a>
Reactivity	Human
Host	Mouse
Clonality	monoclonal
Isotype	IgG1,k
Calculated MW	77847

**RANBP9 Antibody - Additional Information****Gene ID** 10048**Other Names**

Ran-binding protein 9, RanBP9, BPM-L, BPM90, Ran-binding protein M, RanBPM, RanBP7, RANBP9, RANBPM

**Target/Specificity**

This RANBP9 antibody is generated from a mouse immunized with a recombinant protein between 1-388 amino acids from the human RANBP9.

**Dilution**

WB~~1:2000-1:4000

IHC-P~~1:25

E~~Use at an assay dependent concentration.

**Format**

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

RANBP9 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**RANBP9 Antibody - Protein Information****Name** RANBP9**Synonyms** RANBPM

**Function** May act as scaffolding protein, and as adapter protein to couple membrane receptors to intracellular signaling pathways (Probable). Acts as a mediator of cell spreading and actin cytoskeleton rearrangement (PubMed:[18710924](#)). Core component of the CTLH E3 ubiquitin-protein ligase complex that selectively accepts ubiquitin from UBE2H and mediates ubiquitination and subsequent proteasomal degradation of the transcription factor HBP1 (PubMed:[29911972](#)). May be involved in signaling of ITGB2/LFA-1 and other integrins (PubMed:[14722085](#)). Enhances HGF-MET signaling by recruiting Sos and activating the Ras pathway (PubMed:[12147692](#)). Enhances dihydrotestosterone-induced transactivation activity of AR, as well as dexamethasone-induced transactivation activity of NR3C1, but not affect estrogen-induced transactivation (PubMed:[12361945](#), PubMed:[18222118](#)). Stabilizes TP73 isoform Alpha, probably by inhibiting its ubiquitination, and increases its proapoptotic activity (PubMed:[15558019](#)). Inhibits the kinase activity of DYRK1A and DYRK1B. Inhibits FMR1 binding to RNA.

#### **Cellular Location**

Cytoplasm. Nucleus. Cell membrane; Peripheral membrane protein. Note=The unphosphorylated form is predominantly cytoplasmic. A phosphorylated form is associated with the plasma membrane.

#### **Tissue Location**

Ubiquitously expressed, with highest levels in testes, placenta, heart, and muscle, and lowest levels in lung. Within the brain, expressed predominantly by neurons in the gray matter of cortex, the granular layer of cerebellum and the Purkinje cells


### **RANBP9 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **RANBP9 Antibody - Images**





All lanes : Anti-RANBP9 Antibody at 1:2000-1:4000 dilution Lane 1: HeLa whole cell lysate Lane 2: U-2OS whole cell lysate Lane 3: Jurkat whole cell lysate Lane 4: CCRF-CEM whole cell lysate Lane 5: MDA-MB-231 whole cell lysate Lane 6: MOLT-4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 78 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AM8615b staining RANBP9 in human heart tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 3% BSA for 0.5 hour at room temperature; antigen retrieval was by heat mediation with a citrate buffer (pH6). Samples were incubated with primary antibody (1/25) for 1 hour at 37°C. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.

### **RANBP9 Antibody - Background**

May act as an adapter protein to couple membrane receptors to intracellular signaling pathways. May be involved in signaling of ITGB2/LFA-1 and other integrins. Enhances HGF-MET signaling by recruiting Sos and activating the Ras pathway. Enhances dihydrotestosterone-induced transactivation activity of AR, as well as dexamethasone-induced transactivation activity of NR3C1, but not affect estrogen-induced transactivation. Stabilizes TP73 isoform Alpha, probably by inhibiting its ubiquitination, and increases its proapoptotic activity. Inhibits the kinase activity of DYRK1A and DYRK1B. Inhibits FMR1 binding to RNA (By similarity).

**RANBP9 Antibody - References**

- Nishitani H.,et al.Gene 272:25-33(2001).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Mungall A.J.,et al.Nature 425:805-811(2003).  
Nakamura M.,et al.J. Cell Biol. 143:1041-1052(1998).  
Wang Y.,et al.Biochem. Biophys. Res. Commun. 297:148-153(2002).