

GDF15 Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AM8725b

Specification

GDF15 Antibody - Product Information

Application WB,E **Primary Accession** 099988 Reactivity Human Predicted Human Host Mouse Clonality monoclonal Isotype IgG1, ĸ Calculated MW 34140

GDF15 Antibody - Additional Information

Gene ID 9518

Other Names

Growth/differentiation factor 15, GDF-15, Macrophage inhibitory cytokine 1, MIC-1, NSAID-activated gene 1 protein, NAG-1, NSAID-regulated gene 1 protein, NRG-1, Placental TGF-beta, Placental bone morphogenetic protein, Prostate differentiation factor, GDF15 (HGNC:30142)

Target/Specificity

This GDF15 antibody is generated from a mouse immunized with a recombinant protein from the human GDF15.

Dilution

WB~~1:4000

Format

Purified monoclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein G column, followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GDF15 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

GDF15 Antibody - Protein Information

Name GDF15 (<u>HGNC:30142</u>)



Function Regulates food intake, energy expenditure and body weight in response to metabolic and toxin-induced stresses (PubMed:28953886, PubMed:28846097, PubMed:28846098, PubMed:28846099, PubMed:23468844, PubMed:29046435). Binds to its receptor, GFRAL, and activates GFRAL- expressing neurons localized in the area postrema and nucleus tractus solitarius of the brainstem (PubMed:28953886, PubMed:28846097, PubMed:28846098, PubMed:28846099). It then triggers the activation of neurons localized within the parabrachial nucleus and central amygdala, which constitutes part of the 'emergency circuit' that shapes feeding responses to stressful conditions (PubMed:28953886). On hepatocytes, inhibits growth hormone signaling (By similarity).

Cellular Location Secreted

Tissue Location


Highly expressed in placenta, with lower levels in prostate and colon and some expression in kidney (PubMed:9348093) Detected in plasma (at protein level) (PubMed:28572090, PubMed:29046435).

GDF15 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

GDF15 Antibody - Images



Anti-GDF15 Antibody at 1:4000 dilution + HT-1080 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-mouse IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 34 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

GDF15 Antibody - Background





Tel: 858.875.1900 Fax: 858.875.1999

Regulates food intake, energy expenditure and body weight in response to metabolic and toxin-induced stresses (PubMed:28953886, PubMed:28846097, PubMed:28846098, PubMed:28846099, PubMed:23468844, PubMed:29046435). Binds to its receptor, GFRAL, and activates GFRAL-expressing neurons localized in the area postrema and nucleus tractus solitarius of the brainstem (PubMed:28953886, PubMed:28846097, PubMed:28846098, PubMed:28846099). It then triggers the activation of neurons localized within the parabrachial nucleus and central amygdala, which contitutes part of the 'emergency circuit' that shapes feeding responses to stressful conditions (PubMed:28953886). On hepatocytes, inhibits growth hormone signaling (By similarity).

GDF15 Antibody - References

Hromas R., et al. Biochim. Biophys. Acta 1354:40-44(1997). Yokoyama-Kobayashi M., et al.J. Biochem. 122:622-626(1997). Bootcov M.R., et al. Proc. Natl. Acad. Sci. U.S.A. 94:11514-11519(1997). Paralkar V.M., et al.J. Biol. Chem. 273:13760-13767(1998). Kalnine N., et al. Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.