

Phospho-Thr286 CaM Kinase II Antibody
Affinity purified rabbit polyclonal antibody
Catalog # AN1001**Specification**

Phospho-Thr286 CaM Kinase II Antibody - Product Information

Application	WB
Primary Accession	P11275
Reactivity	Rat
Predicted	Human, Mouse, Xenopus
Host	Rabbit
Clonality	polyclonal
Calculated MW	50/60 KDa

Phospho-Thr286 CaM Kinase II Antibody - Additional Information

Gene ID	25400
Gene Name	CAMK2A/B

Other Names

Calcium/calmodulin-dependent protein kinase type II subunit alpha, CaM kinase II subunit alpha, CaMK-II subunit alpha, Camk2a

Target/Specificity

Synthetic phospho-peptide corresponding to amino acid residues surrounding Thr286 conjugated to KLH.

Dilution

WB~~ 1:1000

Format

Prepared from rabbit serum by affinity purification via sequential chromatography on phospho- and dephosphopeptide affinity columns.

Antibody Specificity

Specific for the ~50k α -CaM Kinase II and the ~60k β -CaM Kinase II proteins phosphorylated at Thr286. Immunolabeling is blocked by the λ -phosphatase treatment.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Phospho-Thr286 CaM Kinase II Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Phospho-Thr286 CaM Kinase II Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-Thr286 CaM Kinase II Antibody - Images



Western blot of rat brain lysate showing specific immunolabeling of the ~50k α - and the ~60k β -CaM Kinase II phosphorylated at Thr286 (Control). The phosphospecificity of this labeling is shown in the second lane (lambda-phosphatase: λ -Ptase). The blot is identical to the control except that it was incubated in λ -Ptase (1200 units for 30 min) before being exposed to the Anti-Thr286 CaM Kinase II. The immunolabeling is completely eliminated by treatment with λ -Ptase.

Phospho-Thr286 CaM Kinase II Antibody - Background

Ca²⁺/calmodulin-dependent protein kinase II (CaM Kinase II) is a multi-functional calcium and calmodulin-dependent protein kinase that mediates cellular responses to a wide variety of intercellular signals (Kennedy, 1998; Schulman and Hanson, 1993). CaM Kinase II has been shown to regulate diverse cellular functions including synaptic plasticity, neurotransmitter synthesis and release, gene expression, ion channel function, carbohydrate metabolism, cytoskeletal function, and Ca²⁺-homeostasis (Gleason et al., 2003; Soderling, 2000; Hudmon and Schulman, 2002). Phosphorylation of Thr286 on the kinase produces an autonomously active form of CaM Kinase II (Meng et al., 2003; Picciotto et al., 1993). Autophosphorylation of Thr305 inhibits the activity CaM Kinase II. Phosphorylation at this site appears to reduce the association of CaM Kinase II with the PSD and reduce LTP and learning (Elgersma et al., 2002).

Phospho-Thr286 CaM Kinase II Antibody - References

Elgersma Y, Fedorov NB, Ikonen S, Choi ES, Elgersma M, Carvalho OM, Giese KP, Silva AJ (2002) Inhibitory autophosphorylation of CaMKII controls PSD association, plasticity, and learning. *Neuron* 36:493-505.

Gleason MR, Higashijima S, Dallman J, Liu K, Mandel G, Fetcho JR (2003) Translocation of CaM

kinase II to synaptic sites in vivo. *Nature Neurosci* 6:217-218.

Hudmon A, Schulman H (2002) Neuronal Ca²⁺/calmodulin-dependent protein kinase II: The role of structure and autoregulation in cellular function. *Annu Rev Biochem* 71:473-510.

Kennedy MB (1998) Signal transduction molecules at the glutamatergic postsynaptic membrane. *Brain Res Rev* 26:243-257.

Meng FJ, Guo J, Zhang QG, Song B, Zhang GY (2003) Autophosphorylated calcium/calmodulin-dependent protein kinase IIa (CaMKIIa) reversibly targets to and phosphorylates N-methyl-D-aspartate receptor subunit 2B (NR2B) in cerebral ischemia and reperfusion in hippocampus of rats. *Brain Res* 967:161-169.

Picciotto MR, Czernik AJ, Nairn AC (1993) Calcium/calmodulin-dependent protein kinase I. cDNA cloning and identification of autophosphorylation site. *J Biol Chem* 268:26512-26521.

Schulman H, Hanson PI (1993) Multifunctional Ca²⁺/calmodulin-dependent protein kinase. *Neurochem Res* 18:65-77.

Soderling TR (2000) CaM-kinases: modulators of synaptic plasticity. *Curr Opin Neurobiol* 10:375-380.

Sun X, Milovanovic M, Zhao Y, Wolf ME. (2008) Acute and chronic dopamine receptor stimulation modulates AMPA receptor trafficking in nucleus accumbens neurons cocultured with prefrontal cortex neurons. *J Neurosci.* Apr 16;28(16):4216-30.

Rebekah S. Vest, Heather O'Leary, Steven J. Coultrap, Mark S. Kindy, and K. Ulrich Bayer (2010) Effective Post-insult Neuroprotection by a Novel Ca²⁺/ Calmodulin-dependent Protein Kinase II (CaMKII) Inhibitor. *J. Biol. Chem.*, 285: 20675 - 20682.

Kurtis D. Davies, Susan M. Goebel-Goody, Steven J. Coultrap, and Michael D. Browning (2008) Long Term Synaptic Depression That Is Associated with GluR1 Dephosphorylation but Not -Amino-3-hydroxy-5-methyl-4-isoxazolepropionic Acid (AMPA) Receptor Internalization *J. Biol. Chem.*, 283: 33138 - 33146.

Xiu Sun, Michael Milovanovic, Yun Zhao, and Marina E. Wolf (2008) Acute and Chronic Dopamine Stimulation Modulates AMPA Receptor Trafficking in Nucleus Accumbens Neurons Cocultured with Prefrontal Cortex Neurons. *J. Neurosci.*, 28: 4216 - 4230

Rebekah S. Vest, Kurtis D. Davies, Heather O'Leary, J. David Port, and K. Ulrich Bayer (2007) Dual Mechanism of a Natural CaMKII Inhibitor. *Mol. Biol. Cell*, 18: 5024 - 5033.

Matthew Townsend, Tapan Mehta, and Dennis J. Selkoe (2007) Soluble A Inhibits Specific Signal Transduction Cascades Common to the Insulin Receptor Pathway. *J. Biol. Chem.*, 282: 33305 - 33312