

GABAA Receptor, β2-Subunit Antibody

Affinity purified rabbit polyclonal antibody Catalog # AN1042

Specification

GABAA Receptor, β2-Subunit Antibody - Product Information

Application Primary Accession Reactivity Predicted Host Clonality Calculated MW WB <u>P63138</u> Rat Human, Mouse, Monkey Rabbit polyclonal 55 KDa

GABAA Receptor, β2-Subunit Antibody - Additional Information

Gene ID25451Gene NameGABRB2Other NamesGabraGamma-aminobutyric acid receptor subunit beta-2, GABA(A) receptor subunit beta-2, Gabrb2,
Gabrb-2

Target/Specificity Fusion protein from the cytoplasmic loop of the beta 2 subunit.

Dilution WB~~ 1:1000

Format Prepared from rabbit serum by affinity purification using a column to which the fusion protein immunogen was coupled.

Antibody Specificity Specific for the \sim 55k β 2-subunit of the GABAA receptor in Western blots.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GABAA Receptor, β 2-Subunit Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice

GABAA Receptor, β2-Subunit Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

GABAA Receptor, β2-Subunit Antibody - Images



Western blot of 7 μ g of rat cerebellum (Cb) showing specificimmunolabeling of the ~55k β 2-subunit of the GABAA-R.

GABAA Receptor, β2-Subunit Antibody - Background

Gamma-aminobutyric acid (GABA) is the primary inhibitory neurotransmitter in the central nervous system, causing a hyperpolarization of the membrane through the opening of a Cl– channel associated with the GABAA receptor (GABAA-R) subtype. GABAA-Rs are important therapeutic targets for a range of sedative, anxiolytic, and hypnotic agents and are implicated in several diseases including epilepsy, anxiety, depression, and substance abuse. The GABAA-R is a multimeric subunit complex. To date six α s, four β s and four γ s, plus alternative splicing variants of some of these subunits, have been identified (Olsen and Tobin, 1990; Whiting et al., 1999; Ogris et al., 2004). Injection in oocytes or mammalian cell lines of cRNA coding for α - and β -subunits results in the expression of functional GABAA-Rs sensitive to GABA. However, coexpression of a γ -subunit is required for benzodiazepine modulation. The various effects of the benzodiazepines in brain may also be mediated via different α -subunits of the receptor (McKernan et al., 2000; Mehta and Ticku, 1998; Ogris et al., 2004; Pöltl et al., 2003).

GABAA Receptor, β2-Subunit Antibody - References

McKernan RM, et al. (2000) Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABAA receptor α 1-subtype. Nature Neurosci 3:587-592.

Mehta AK, Ticku MK (1998) Prevalence of the GABAA receptor assemblies containing α 1-subunit in the rat cerebellum and cerebral cortex as determined by immunoprecipitation: Lack of modulation by chronic ethanol administration. Mol Brain Res 67:194-199.

Ogris W, Pöltl A, Hauer B, Ernst M, Oberto A, Wulff P, Höger H, Wisden W, Sieghart W (2004) Affinity of various benzodiazepine site ligands in mice with a point mutation in the GABAA receptor γ 2-subunit. Biochem Pharmacol 68:1621-1629.



Olsen RW, Tobin AJ (1990) Molecular biology of GABAA receptors. FASEB 4:1469-1480. Pöltl A, Hauer B, Fuchs K, Tretter V, Sieghart W (2003) Subunit composition and quantitative importance of GABAA receptor subtypes in the cerebellum of mouse and rat. J Neurochem 87:1444-1455.

Whiting PJ, Bonnert TP, McKernan RM, Farrar S, Le Bourdellès B, Heavens RP, Smith DW, Hewson L, Rigby MR, Sirinathsinghji DJS, Thompson SA, Wafford KA (1999) Molecular and functional diversity of the expanding GABAA receptor gene family. Ann NY Acad Sci 868:645-653.