

NMDA Receptor, NR1 Subunit, N1 Splice Variant Insert Antibody
Affinity purified rabbit polyclonal antibody
Catalog # AN1051**Specification**

NMDA Receptor, NR1 Subunit, N1 Splice Variant Insert Antibody - Product Information

Application	WB
Primary Accession	P35439
Reactivity	Mouse, Rat
Predicted	Chicken, Human, Monkey, Xenopus
Host	Rabbit
Clonality	polyclonal
Calculated MW	120 KDa

NMDA Receptor, NR1 Subunit, N1 Splice Variant Insert Antibody - Additional Information

Gene ID	24408
Gene Name	GRIN1

Other Names

Glutamate receptor ionotropic, NMDA 1, GluN1, Glutamate [NMDA] receptor subunit zeta-1, N-methyl-D-aspartate receptor subunit NR1, NMD-R1, Grin1, Nmdar1

Target/Specificity

Synthetic peptide corresponding to amino acid residues specific to the NR1 subunit, N1 splice variant insert conjugated to KLH.

Dilution

WB~~ 1:1000

Format

Prepared from rabbit serum by affinity purification using a column to which the peptide immunogen was coupled.

Antibody Specificity

Specific for the ~120k NR1 subunit, N1 splice variant insert protein of the NMDA receptor. Does not recognize NR1 subunits of the NMDA receptor that do not contain the N1 insert.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NMDA Receptor, NR1 Subunit, N1 Splice Variant Insert Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

NMDA Receptor, NR1 Subunit, N1 Splice Variant Insert Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NMDA Receptor, NR1 Subunit, N1 Splice Variant Insert Antibody - Images



Western blot of 10 ug of HEK 293 cells expressing: Lane 1 - NR1 subunit containing the N1 and C2' Insert showing specific immunolabeling of the ~120k NR1 subunit of the NMDA receptor containing the N1 splice variant insert.; Lane 2 - NR1 subunit containing only the C2 Insert; Lane 3 - NR1 subunit containing the C1 and C2' Insert; Lane 4 - without NR1 expression (Mock).

NMDA Receptor, NR1 Subunit, N1 Splice Variant Insert Antibody - Background

The ion channels activated by glutamate that are sensitive to N-methyl-D-aspartate (NMDA) are designated NMDA Receptors (NMDAR). The NMDAR plays an essential role in memory, neuronal development and it has also been implicated in several disorders of the central nervous system including Alzheimer's, epilepsy and ischemic neuronal cell death (Grosshans et al., 2002; Wenthold et al., 2003; Carroll and Zukin, 2002). There are a number of different splice variants of the NR1-Subunit (Foldes et al., 1994; Zukin and Bennett, 1995). Differential splicing of three exons in the NR1-Subunit generates up to eight NR1-Subunit splice variants and 7 of these have been identified in cDNA libraries. These exons encode a 21 amino acid N-terminal domain (N1) and adjacent sequences in the C-terminus (C1 and C2). Splicing out the C2 cassette eliminates the first stop codon and produces a new reading frame that generates a new sequence of 22 amino acids (C2'). Considerable attention has been focused on the distribution and expression of these splice variants that may affect the functional properties and regulation of the NMDAR.

NMDA Receptor, NR1 Subunit, N1 Splice Variant Insert Antibody - References

- Carroll RC, Zukin RS (2002) NMDA-receptor trafficking and targeting: implications for synaptic transmission and plasticity. *Trends Neurosci* 25:571-577.
- Foldes RL, Rampersad V, Kamboj RK (1994) Cloning and sequence analysis of additional splice variants encoding human N-methyl-D-aspartate receptor (hNR1) subunits. *Gene* 147:303-304.
- Grosshans DR, Clayton DA, Coultrap SJ, Browning MD (2002) LTP leads to rapid surface expression of NMDA but not AMPA receptors in adult rat CA1. *Nat Neurosci* 5:27-33.
- Wenthold RJ, Prybylowski K, Standley S, Sans N, Petralia RS (2003) Trafficking of NMDA receptors. *Annu Rev Pharmacol Toxicol* 43:335-358.
- Zukin RS, Bennett MVL (1995) Alternatively spliced isoforms of the NMDAR1 receptor subunit.

Trends Neurosci 18:306-313.

Note: Dr. Michael Browning, a co-author of one of the cited papers, is President and founder of PhosphoSolutions