

Dopamine Transporter, C-Terminus Antibody
Affinity purified rabbit polyclonal antibody
Catalog # AN1068**Specification**

Dopamine Transporter, C-Terminus Antibody - Product Information

Application	WB
Primary Accession	Q01959
Reactivity	Human, Mouse, Monkey
Host	rabbit
Clonality	polyclonal
Calculated MW	88 KDa

Dopamine Transporter, C-Terminus Antibody - Additional Information

Gene Name **SLC6A3**

Other Names

Sodium-dependent dopamine transporter, DA transporter, DAT, Solute carrier family 6 member 3, SLC6A3, DAT1

Target/Specificity

Peptide from the intracellular C-terminus region of human dopamine transporter (DAT), conjugated to keyhole limpet hemocyanin (KLH).

Dilution

WB~~ 1:1000

Format

Prepared from rabbit serum by affinity purification using a SulfoLink® column matrix to which the peptide immunogen was coupled.

Antibody Specificity

Specific for the ~88k DAT protein in Western blots.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Dopamine Transporter, C-Terminus Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping


Blue Ice

Dopamine Transporter, C-Terminus Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Dopamine Transporter, C-Terminus Antibody - Images



Western blot of human caudate lysate showing specific immunolabeling of the ~88k DAT protein.

Dopamine Transporter, C-Terminus Antibody - Background

The dopamine transporter (DAT) is responsible for the reaccumulation of dopamine after it has been released. DAT antibodies and antibodies for other markers of catecholamine biosynthesis are widely used as markers for dopaminergic and noradrenergic neurons in a variety of applications including depression, schizophrenia, Parkinson's disease and drug abuse (Kish et al., 2001; Zhu et al., 2000; Zhu et al., 1999). Levels of DAT protein expression are altered by chronic drug administration (Wilson et al., 1996).

Dopamine Transporter, C-Terminus Antibody - References

- Kish SJ, Kalasinsky KS, Derkach P, Schmunk GA, Guttman M, Ang L, Adams V, Furukawa Y, Haycock JW (2001) Striatal dopaminergic and serotonergic markers in human heroin users. *Neuropsychopharmacology* 24:561-567.
- Wilson JM, Kalasinsky KS, Levey AI, Bergeron C, Reiber G, Anthony RM, Schmunk GA, Shannak K, Haycock JW, Kish SJ (1996) Striatal dopamine nerve terminal markers in human, chronic methamphetamine users. *Nat Med* 2:699-703.
- Zhu MY, Klimek V, Haycock JW, Ordway GA (2000) Quantitation of tyrosine hydroxylase protein in the locus coeruleus from postmortem human brain. *J Neurosci Meth* 99:37-44.
- Zhu MY, Klimek V, Dilley GE, Haycock JW, Stockmeier C, Overholser JC, Meltzer HY, Ordway GA (1999) Elevated levels of tyrosine hydroxylase in the locus coeruleus in major depression. *Biol Psychiatry* 46:1275-1286.