

Neurofilament M (NF-M) Antibody

Mouse monoclonal antibody Catalog # AN1148

Specification

Neurofilament M (NF-M) Antibody - Product Information

Application Primary Accession Reactivity Predicted Host Clonality Isotype Calculated MW IF, WB <u>P12839</u> Rat Chicken, Human, Mouse Mouse monoclonal IgG1 145 KDa

Neurofilament M (NF-M) Antibody - Additional Information

Gene ID24588Gene NameNEFMOther NamesNeurofilament medium polypeptide, NF-M, 160 kDa neurofilament protein, Neurofilament 3,
Neurofilament triplet M protein, Nefm, Nef3, Nfm

Target/Specificity Preparation containing the extreme C-terminus expressed in and purified from E. Coli.

Dilution IF~~ 1:100 WB~~ 1:1000

Format Unpurified, concentrated culture supernatant.

Antibody Specificity Specific for the ~145k neurofilament M protein.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Neurofilament M (NF-M) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice


Neurofilament M (NF-M) Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Neurofilament M (NF-M) Antibody - Images



Western blot of rat cortex lysate showing specific immunolabeling of the ~ 145k NF-M protein.



Immunostaining of cultured rat neurons showing labeling of NF-M (green) in mature neurons. Neurofilament M (NF-M) Antibody - Background

Neurofilaments are the 10nm or intermediate filament proteins found specifically in neurons, and are composed predominantly of three major proteins called NF-L, NF-M and NF-H (1). NF-M is the neurofilament middle or medium molecular weight polypeptide and runs on SDS-PAGE gels at 145-160 kDa, with some variability across species boundaries. Antibodies to NF-M are useful for identifying neuronal cells and their processes in tissue sections and in tissue culture. NF-M antibodies can also be useful to visualize neurofilament accumulations seen in many neurological



diseases, such as Amyotrophic Lateral Sclerosis (Lou Gehrig's disease) and Alzheimer's disease (2).

Neurofilament M (NF-M) Antibody - References

1.

Harris, J., Ayyub, C. and Shaw G. (1991) A molecular dissection of the carboxyterminal tails of the major neurofilament subunits NF-M and NF-H. J Neurosci Res 30:47-62.

2.

Hu YY, He SS, Wang XC, Duan QH, Khatoon S, Igbal K, Grundke-Igbal I, Wang JZ (2002) Elevated levels of phosphorylated neurofilament proteins in cerebrospinal fluid of Alzheimer disease patients. Neurosci Lett 320(3):156-60.