

**Anti-cdc2 (Tyr15) Antibody**

**Our Anti-cdc2 (Tyr15) rabbit polyclonal phosphospecific primary antibody from PhosphoSolutions is pr  
Catalog # AN1330**

**Specification**

---

**Anti-cdc2 (Tyr15) Antibody - Product Information**

Primary Accession	<a href="#">P39951</a>
Reactivity	<b>Bovine</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>IgG</b>
Calculated MW	<b>34135</b>

**Anti-cdc2 (Tyr15) Antibody - Additional Information**

Gene ID **54237**

**Other Names**

Cdc 2 antibody, Cdc2 antibody, CDC28A antibody, CDK-1 antibody, CDK 1 antibody, CDK1 antibody, CDK1\_HUMAN antibody, CDKN1 antibody, CELL CYCLE CONTROLLER CDC2 antibody, Cell division control protein 2 antibody, Cell division control protein 2 homolog antibody, Cell division cycle 2 G1 to S and G2 to M antibody, Cell division protein kinase 1 antibody, Cell Division Cycle 2 Protein antibody, Cyclin Dependent Kinase 1 antibody, Cyclin-dependent kinase 1 antibody, DKFZp686L20222 antibody, MGC111195 antibody, p34 Cdk1 antibody, p34 protein kinase antibody, P34CDC2 antibody

**Target/Specificity**

Cdc2 is a highly conserved protein serine kinase that plays a key role in regulation of the cell cycle (Maller, 1991). The ability of cdc2 to exercise control over the cell cycle is dependent upon the phosphorylation of Tyr-15 in cdc2 (Nakamizo et al., 2002). cdc2 expression in brain has been linked to both neurogenesis and apoptosis (Konishi and Bonni, 2003; Dranovsky et al., 2001; Okano et al., 1996).

**Format**

Antigen Affinity Purified from Pooled Serum

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Anti-cdc2 (Tyr15) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Shipping**


Blue Ice

**Anti-cdc2 (Tyr15) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-cdc2 (Tyr15) Antibody - Images



Western blot of human T47D cells showing specific immunolabeling of the ~34 kDa cdc2 phosphorylated at Tyr15 (Control, Lane 1). Treatment with EGF (30 ng per ml for 30 min) caused dephosphorylation of the Tyr15 on cdc2 (Lane 2). Phosphospecificity is shown in lanes 3 and 4, where the immunolabeling is completely eliminated by blot treatment with lambda phosphatase, λ-Ptase (1200 units for 30 min).

#### Anti-cdc2 (Tyr15) Antibody - Background

Cdc2 is a highly conserved protein serine kinase that plays a key role in regulation of the cell cycle (Maller, 1991). The ability of cdc2 to exercise control over the cell cycle is dependent upon the phosphorylation of Tyr-15 in cdc2 (Nakamizo et al., 2002). cdc2 expression in brain has been linked to both neurogenesis and apoptosis (Konishi and Bonni, 2003; Dranovsky et al., 2001; Okano et al., 1996).