

Anti-Collagen 1, alpha 1 telopeptide Antibody

Our Anti-Collagen 1, alpha 1 telopeptide primary antibody from PhosphoSolutions is rabbit polyclonal Catalog # AN1341

Specification

Anti-Collagen 1, alpha 1 telopeptide Antibody - Product Information

ApplicationWBPrimary AccessionP02452ReactivityBovine, Chicken, DrosophilaHostRabbitClonalityPolyclonalIsotypeIgGCalculated MW138911

Anti-Collagen 1, alpha 1 telopeptide Antibody - Additional Information

Gene ID

1277

Other Names

Alpha 1 type I collagen antibody, Alpha 2 type I collagen antibody, alpha 2 type I procollagen antibody, alpha 2(I) procollagen antibody, alpha 2(I)-collagen antibody, Alpha-1 type I collagen antibody, alpha1(I) procollagen antibody, CO1A1_HUMAN antibody, COL1A1 antibody, COL1A2 antibody, collagen alpha 1 chain type I antibody, Collagen alpha-1(I) chain antibody, collagen alpha-1(I) chain preproprotein antibody, Collagen I alpha 1 polypeptide antibody, Collagen I alpha 2 polypeptide antibody, collagen of skin tendon and bone, alpha-1 chain antibody, collagen of skin tendon and bone alpha-2 chain antibody, Collagen type I alpha 1 antibody, Collagen type I alpha 2 antibody, EDSC antibody, OI1 antibody, OI2 antibody, OI3 antibody, OI4 antibody, pro-alpha-1 collagen type 1 antibody, type I proalpha 1 antibody, type I procollagen alpha 1 chain antibody, Type I procollagen antibody

Target/Specificity

Collagen is an extracellular matrix protein that serves as a scaffold defining the shape and mechanical properties of many tissues and organs including skin, tendon, artery walls, fibrocartilage, bone and teeth. Type 1 collagen is the most abundant protein in mammals. Collagens are synthesized with N-terminal and C-terminal propeptides that are cleaved during maturation and secretion. After cleavage of the propeptides, the most N-terminal and C-terminal remaining sequences are known as telopeptides. Mutations in the collagen 1, alpha 1 gene (COL1A1) are known to cause osteogenesis imperfecta (aka brittle bone disease) (Byers 1989). Furthermore, mutations found in the first 90 residues of the procollagen N-propeptide leading to a combined osteogenesis imperfecta and Ehlers-Danlos syndrome (EDS) phenotype (Cabral et al., 2005).

Dilution WB~~1:1000

Format Antigen Affinity Purified

Storage



Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Anti-Collagen 1, alpha 1 telopeptide Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice

Anti-Collagen 1, alpha 1 telopeptide Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Collagen 1, alpha 1 telopeptide Antibody - Images



Immunostaining of formaldehyde-fixed fibrotic mouse lung tissue. The antibody recognizes mature collagen I (cat. 322-COLT, 1:100, red) that has formed fibrils in the extracellular matrix. The blue is staining nuclear DNA.





Immunofluorescence of a transversal section from mouse spinal cord labeling fibers of collagen (cat. : 322-COLT, red, 1:200) from the tissue scaffold. The blue is DAPI counterstaining nuclear DNA. Image kindly provided by Dr. Rodolfo Gatto and Dr. Gerardo Morfini, Department of Anatomy and Cell biology, University of Illinois at Chicago.



Western blot of rat lung lysate showing specific immunolabeling of the \sim 140 kDa collagen protein.

Anti-Collagen 1, alpha 1 telopeptide Antibody - Background

Collagen is an extracellular matrix protein that serves as a scaffold defining the shape and mechanical properties of many tissues and organs including skin, tendon, artery walls, fibrocartilage, bone and teeth. Type 1 collagen is the most abundant protein in mammals. Collagens are synthesized with N-terminal and C-terminal propeptides that are cleaved during maturation and secretion. After cleavage of the propeptides, the most N-terminal and C-terminal remaining sequences are known as telopeptides. Mutations in the collagen 1, alpha 1 gene (COL1A1) are known to cause osteogenesis imperfecta (aka brittle bone disease) (Byers 1989). Furthermore, mutations found in the first 90 residues of the procollagen N-propeptide leading to a combined osteogenesis imperfecta and Ehlers-Danlos syndrome (EDS) phenotype (Cabral et al., 2005).