

**Anti-Dynamin (Ser778) Antibody**

**Our Anti-Dynamin (Ser778) sheep polyclonal phosphospecific primary antibody from PhosphoSolutions is  
Catalog # AN1371**

**Specification**

---

**Anti-Dynamin (Ser778) Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Primary Accession | <a href="#">P21575</a> |
| Host              | <b>Sheep</b>           |
| Clonality         | <b>Polyclonal</b>      |
| Isotype           | <b>IgG</b>             |
| Calculated MW     | <b>97295</b>           |

**Anti-Dynamin (Ser778) Antibody - Additional Information**

Gene ID **140694**

**Other Names**

B dynamin antibody, D100 antibody, DNM 1 antibody, DNM antibody, DNM1 antibody, DYN1\_HUMAN antibody, Dynamin antibody, Dynamin-1 antibody, Dynamin1 antibody

**Target/Specificity**

Dynamin is a member of a group of nerve terminal proteins called dephosphins that regulate synaptic vesicle endocytosis (Cousin et al., 2001; Graham et al., 2002; Tsuboi et al., 2002). Cyclin dependent protein kinase 5 phosphorylates dynamin at Ser-774 and Ser-778 that are the phosphorylation sites on dynamin phosphorylated in vivo (Tan et al., 2003). Phosphorylation of these sites on dynamin is thought to play a key role in synaptic vesicle trafficking.

**Format**

Antigen Affinity Purified from Pooled Serum

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Anti-Dynamin (Ser778) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Shipping**

Blue Ice


**Anti-Dynamin (Ser778) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-Dynamin (Ser778) Antibody - Images



Western blot of rat hippocampal lysate stimulated with forskolin showing specific immunolabeling of the ~95 kDa dynamin phosphorylated at Ser778 in the first lane (-). Phosphospecificity is shown in the second lane (+) where immunolabeling is completely eliminated by blot treatment with lambda phosphatase ( $\lambda$ -Ptase, 1200 units for 30 min).

#### Anti-Dynamin (Ser778) Antibody - Background

Dynamin is a member of a group of nerve terminal proteins called dephosphins that regulate synaptic vesicle endocytosis (Cousin et al., 2001; Graham et al., 2002; Tsuboi et al., 2002). Cyclin dependent protein kinase 5 phosphorylates dynamin at Ser-774 and Ser-778 that are the phosphorylation sites on dynamin phosphorylated in vivo (Tan et al., 2003). Phosphorylation of these sites on dynamin is thought to play a key role in synaptic vesicle trafficking.