

**Anti-EphrinB (Tyr298) Antibody**

**Our Anti-EphrinB (Tyr298) rabbit polyclonal phosphospecific primary antibody from PhosphoSolutions i**  
**Catalog # AN1375**

**Specification**

---

**Anti-EphrinB (Tyr298) Antibody - Product Information**

Primary Accession	<a href="#">P28693</a>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>IgG</b>
Calculated MW	<b>111963</b>

**Anti-EphrinB (Tyr298) Antibody - Additional Information**

Gene ID **396513**

**Other Names**

CEK5 ligand antibody, EFNB1 antibody, ELK L antibody, ELK ligand antibody, ELKL antibody, EPH related receptor tyrosine kinase ligand 2 antibody, Ephrin B1 antibody, Ephrin B2 antibody, LERK2 antibody

**Target/Specificity**

EphrinB proteins are thought to play key roles in cellular functions as diverse as neuronal migration and blood vessel development (Flanagan and Vanceraeghen, 1998; Dufour et al., 2003; Oike et al., 2002). EphrinB molecules expressed at the membrane surface bind to the EphB family receptors on target cells during cell to cell contact. This interaction leads to cell signaling in the target cell but also generates a reverse signal in the cell expressing EphrinB on its surface. This reverse signaling event is thought to be critical for vessel maturation and neuronal development. Importantly, tyrosine phosphorylation of EphrinB is thought to be a critical component of this reverse signaling event (Palmer et al., 2002). Recent work suggests that phosphorylation of a specific EphrinB residue (Tyr-298) plays a key role in EphrinB signaling (Kalo, et al., 2001).

**Format**

Antigen Affinity Purified from Pooled Serum

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Anti-EphrinB (Tyr298) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Shipping**


Blue Ice

**Anti-EphrinB (Tyr298) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-EphrinB (Tyr298) Antibody - Images**



Western blot of rat testes lysate showing specific immunolabeling of the ~46 kDa EphrinB phosphorylated at Tyr298 in the first lane (-). Phosphospecificity is shown in the second lane (+) where immunolabeling is completely eliminated by blot treatment with lambda phosphatase ( $\lambda$ -Ptase, 1200 units for 30 min).

#### **Anti-EphrinB (Tyr298) Antibody - Background**

EphrinB proteins are thought to play key roles in cellular functions as diverse as neuronal migration and blood vessel development (Flanagan and Vancerahaegen, 1998; Dufour et al., 2003; Oike et al., 2002). EphrinB molecules expressed at the membrane surface bind to the EphB family receptors on target cells during cell to cell contact. This interaction leads to cell signaling in the target cell but also generates a reverse signal in the cell expressing EphrinB on its surface. This reverse signaling event is thought to be critical for vessel maturation and neuronal development. Importantly, tyrosine phosphorylation of EphrinB is thought to be a critical component of this reverse signaling event (Palmer et al., 2002). Recent work suggests that phosphorylation of a specific EphrinB residue (Tyr-298) plays a key role in EphrinB signaling (Kalo, et al., 2001).