

## **Anti-GABAA Receptor γ2 Antibody**

Our Anti-GABAA Receptor γ2 rabbit polyclonal primary antibody from PhosphoSolutions is produced in-h Catalog # AN1403

### Specification

## Anti-GABAA Receptor y2 Antibody - Product Information

WB
<u>P18508</u>
Rabbit
Polyclonal
lgG
54077

### Anti-GABAA Receptor y2 Antibody - Additional Information

Gene ID

29709

Other Names

CAE 2 antibody, CAE2 antibody, ECA 2 antibody, ECA2 antibody, GABA(A) receptor gamma 2 antibody, GABA(A) receptor subunit gamma 2 antibody, GABA(A) receptor subunit gamma-2 antibody, GABRG 2 antibody, GABRγ2 antibody, Gamma aminobutyric acid (GABA) A receptor gamma 2 antibody, Gamma aminobutyric acid A receptor gamma 2 antibody, Gamma aminobutyric acid receptor gamma 2 subunit antibody, Gamma-aminobutyric acid receptor subunit gamma-2 antibody, GBRG2 HUMAN antibody, GEFSP 3 antibody, GEFSP3 antibody

#### **Target/Specificity**

Gamma-aminobutyric acid (GABA) is the primary inhibitory neurotransmitter in the central nervous system, causing a hyperpolarization of the membrane through the opening of a Cl– channel associated with the GABA-A receptor (GABA-A-R) subtype. GABA-A-Rs are important therapeutic targets for a range of sedative, anxiolytic, and hypnotic agents and are implicated in several diseases including epilepsy, anxiety, depression, and substance abuse. The GABA-A-R is a multimeric subunit complex. To date six  $\alpha$ s, four  $\beta$ s and four  $\gamma$ s, plus alternative splicing variants of some of these subunits, have been identified (Olsen and Tobin, 1990; Whiting et al., 1999; Ogris et al., 2004). Injection in oocytes or mammalian cell lines of cRNA coding for  $\alpha$ - and  $\beta$ -subunits results in the expression of functional GABA-A-Rs sensitive to GABA. However, co-expression of a  $\gamma$ -subunit is required for benzodiazepine modulation. The various effects of the benzodiazepines in brain may also be mediated via different  $\alpha$ - subunits of the receptor (McKernan et al., 2000; Mehta and Ticku, 1998; Ogris et al., 2004; Pöltl et al., 2003). Lastly, phosphorylation of  $\beta$ -subunits of the receptor has been shown to modulate GABA-A-R function (Brandon et al., 2003).

Dilution WB~~1:1000

Format Neat Pooled Serum

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.



Precautions

Anti-GABAA Receptor  $\gamma 2$  Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Shipping Blue Ice

# Anti-GABAA Receptor γ2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## Anti-GABAA Receptor y2 Antibody - Images



Western blot of 10  $\mu$ g of rat hippocampal lysate showing specific immunolabeling of the ~46 kDa  $\gamma$ 2-subunit of the GABAA-R.





Immunostaining of rat cerebellum showing labeling of GABAA  $\gamma$ 2 subunit in fuchsia.

## Anti-GABAA Receptor γ2 Antibody - Background

Gamma-aminobutyric acid (GABA) is the primary inhibitory neurotransmitter in the central nervous system, causing a hyperpolarization of the membrane through the opening of a Cl– channel associated with the GABA-A receptor (GABA-A-R) subtype. GABA-A-Rs are important therapeutic targets for a range of sedative, anxiolytic, and hypnotic agents and are implicated in several diseases including epilepsy, anxiety, depression, and substance abuse. The GABA-A-R is a multimeric subunit complex. To date six  $\alpha$ s, four  $\beta$ s and four  $\gamma$ s, plus alternative splicing variants of some of these subunits, have been identified (Olsen and Tobin, 1990; Whiting et al., 1999; Ogris et al., 2004). Injection in oocytes or mammalian cell lines of cRNA coding for  $\alpha$ - and  $\beta$ -subunits results in the expression of functional GABA-A-Rs sensitive to GABA. However, co-expression of a  $\gamma$ -subunit is required for benzodiazepine modulation. The various effects of the benzodiazepines in brain may also be mediated via different  $\alpha$ - subunits of the receptor (McKernan et al., 2000; Mehta and Ticku, 1998; Ogris et al., 2004; Pöltl et al., 2003). Lastly, phosphorylation of  $\beta$ -subunits of the receptor has been shown to modulate GABA-A-R function (Brandon et al., 2003).